



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/67, 15/90, 15/85, 15/62, 5/06, C07K 14/505, 14/61, A61K 48/00		A1	(11) International Publication Number: WO 95/31560 (43) International Publication Date: 23 November 1995 (23.11.95)
(21) International Application Number: PCT/US95/06045 (22) International Filing Date: 11 May 1995 (11.05.95) (30) Priority Data: 08/243,391 13 May 1994 (13.05.94) US (71) Applicant (for all designated States except US): TRAN- SKARYOTIC THERAPIES, INC. [US/US]; 195 Albany Street, Cambridge, MA 02139 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TRECO, Douglas, A. [US/US]; 87 Brantwood Road, Arlington, MA 02174 (US). HEARTLEIN, Michael, W. [US/US]; 167 Reed Farm Road, Boxborough, MA 01719 (US). SELDEN, Richard, F. [US/US]; 106 Bristol Road, Wellesley, MA 02181 (US). (74) Agents: GRANAHAN, Patricia et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US).			(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>
(54) Title: DNA CONSTRUCT FOR EFFECTING HOMOLOGOUS RECOMBINATION AND USES THEREOF			
(57) Abstract <p>The invention relates to constructs comprising: a) a targeting sequence; b) a regulatory sequence; c) an exon; and d) an unpaired splice-donor site. The invention further relates to a method of producing protein <i>in vitro</i> or <i>in vivo</i> comprising the homologous recombination of a construct as described above within a cell. The homologously recombinant cell is then maintained under conditions which will permit transcription and translation, resulting in protein expression. The present invention further relates to homologously recombinant cells, including primary, secondary, or immortalized vertebrate cells, methods of making the cells, methods of homologous recombination to produce fusion genes, methods of altering gene expression in the cells, and methods of making a protein in a cell employing the constructs of the invention.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DNA construct for effecting homologous recombination and uses thereof

Background of the Invention

Current approaches to treating disease by administering therapeutic proteins include in vitro production of therapeutic proteins for conventional pharmaceutical delivery (e.g. intravenous, subcutaneous, or intramuscular injection) and, more recently, gene therapy.

Proteins of therapeutic interest are generally produced by introducing exogenous DNA encoding the protein of therapeutic interest into appropriate cells. For example, exogenous DNA encoding a desired therapeutic protein is introduced into cells, such as immortalized cells in a vector, such as a plasmid, from which the encoded protein is expressed. Further, it has been suggested that endogenous cellular genes and their expression may be modified by gene targeting. See for example, U.S. Patent No. 5,272,071, WO 91/06666, WO 91/06667 and WO 90/11354.

-2-

Presently-available approaches to gene therapy make use of infectious vectors, such as retroviral vectors, which include the genetic material to be expressed. Such approaches have limitations, such as the potential of
5 generating replication-competent virus during vector production; recombination between the therapeutic virus and endogenous retroviral genomes, potentially generating infectious agents with novel cell specificities, host ranges, or increased virulence and cytotoxicity; indepen-
10 dent integration into large numbers of cells, increasing the risk of a tumorigenic insertional event; limited cloning capacity in the retrovirus (which restricts therapeutic applicability) and short-lived in vivo expression of the product of interest. A better approach to provid-
15 ing gene products, particularly one which avoids the limitations and risks associated with presently available methods, would be valuable.

Summary of the Invention

The present invention relates to improved methods for
20 both the in vitro production of therapeutic proteins and for the production and delivery of therapeutic proteins by gene therapy. In the present method, expression of a desired targeted gene in a cell (i.e., a desired endogenous cellular gene) is altered by the introduction, by
25 homologous recombination into the cellular genome at a preselected site, of DNA which includes at least a regulatory sequence, an exon and a splice donor site. These components are introduced into the chromosomal (genomic) DNA in such a manner that this, in effect, results in
30 production of a new transcription unit (in which the regulatory sequence, the exon and the splice donor site present in the DNA construct are operatively linked to the endogenous gene). As a result of introduction of these

-3-

components into the chromosomal DNA, the expression of the desired endogenous gene is altered.

Altered gene expression, as used herein, encompasses activating (or causing to be expressed) a gene which is normally silent (unexpressed) in the cell as obtained, increasing expression of a gene which is not expressed at physiologically significant levels in the cell as obtained, changing the pattern of regulation or induction such that it is different than occurs in the cell as obtained, and reducing (including eliminating) expression of a gene which is expressed in the cell as obtained.

The present invention further relates to DNA constructs useful in the method of altering expression of a target gene. The DNA constructs comprise: (a) a targeting sequence; (b) a regulatory sequence; (c) an exon; and (d) an unpaired splice-donor site. The targeting sequence in the DNA construct directs the integration of elements (a) - (d) into a target gene in a cell such that the elements (b) - (d) are operatively linked to sequences of the endogenous target gene. In another embodiment, the DNA constructs comprise: (a) a targeting sequence, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a) - (f) such that the elements of (b) - (f) are operatively linked to the endogenous gene. The targeting sequence is homologous to the preselected site in the cellular chromosomal DNA with which homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

The following serves to illustrate two embodiments of the present invention, in which the sequences upstream of the human erythropoietin h(EPO) gene are altered to allow expression of hEPO in primary, secondary, or immortalized

-4-

cells which do not express EPO in detectable quantities in their untransfected state as obtained. In embodiment 1, the targeting construct contains two targeting sequences. The first targeting sequence is homologous to sequences 5' of the second targeting sequence, and both sequences are upstream of the hEPO coding region. The targeting construct also contains a regulatory region (the mMT-1 promoter) an exon (human growth hormone (hGH) exon 1) and an unpaired splice-donor site. The product of homologous recombination with this targeting construct is shown in Figure 1.

In embodiment 2, the targeting construct also contains two targeting sequences. The first targeting sequence is homologous to sequences within the endogenous hEPO regulatory region, and the second targeting sequence is homologous to hEPO intron 1. The targeting construct also contains a regulatory region (the mMT-1 promoter), an exon (hGH exon 1) and an unpaired splice-donor site. The product of homologous recombination with this targeting construct is shown in Figure 2.

In these two embodiments, the products of the targeting events are chimeric transcription units which generate a mature mRNA in which the first exon of the hGH gene is positioned upstream of hEPO exons 2-5. The product of transcription, splicing, and translation is a protein in which amino acids 1-4 of the hEPO signal peptide are replaced with amino acid residues 1-3 of hGH. The two embodiments differ with respect to both the relative positions of the regulatory sequences of the targeting construct that are inserted and the specific pattern of splicing that needs to occur to produce the final, processed transcript.

The invention further relates to a method of producing protein in vitro or in vivo through introduction of a construct as described above into host cell chromosomal

-5-

DNA by homologous recombination to produce a homologously recombinant cell. The homologously recombinant cell is then maintained under conditions which will permit transcription, translation and secretion, resulting in production of the protein of interest.

The present invention relates to transfected cells, such as transfected primary or secondary cells (i.e., non-immortalized cells) and transfected immortalized cells, useful for producing proteins, particularly therapeutic proteins, methods of making such cells, methods of using the cells for in vitro protein production, and methods of gene therapy. Cells of the present invention are of vertebrate origin, particularly of mammalian origin, and even more particularly of human origin. Cells produced by the method of the present invention contain DNA which encodes a therapeutic product, DNA which is itself a therapeutic product and/or DNA which causes the transfected cells to express a gene at a higher level or with a pattern of regulation or induction that is different than occurs in the corresponding nontransfected cell.

The present invention also relates to methods by which cells, such as primary, secondary, and immortalized cells, are transfected to include exogenous genetic material, methods of producing clonal cell strains or heterogeneous cell strains, and methods of immunizing animals or producing antibodies in immunized animals, using the transfected primary, secondary, or immortalized cells.

The present invention relates particularly to a method of gene targeting or homologous recombination in eukaryotic cells, such as cells of fungal, plant or animal, e.g., vertebrate, particularly mammalian, and even more particularly, human origin. That is, it relates to a method of introducing DNA into primary, secondary, or immortalized cells of vertebrate origin through homologous recombination, such that the DNA is introduced into genom-

-6-

ic DNA of the primary, secondary, or immortalized cells at a preselected site. The targeting sequences used are selected with reference to the site into which the DNA in the targeting DNA construct is to be inserted. The present invention further relates to homologously recombinant primary, secondary, or immortalized cells, referred to as homologously recombinant (HR) primary, secondary or immortalized cells, produced by the present method and to uses of the HR primary, secondary, or immortalized cells.

10 In one embodiment of the present invention in which expression of a gene is altered, the gene is activated. That is, a gene present in primary, secondary, or immortalized cells of vertebrate origin, which is normally not expressed in the cells as obtained, is activated and, as a result, the encoded protein is expressed. In this embodiment, homologous recombination is used to replace, disable, or disrupt the regulatory region normally associated with the gene in cells as obtained through the insertion of a regulatory sequence which causes the gene to be expressed at levels higher than evident in the corresponding nontransfected cell.

In one embodiment, the activated gene can be further amplified by the inclusion of an amplifiable selectable marker gene which has the property that cells containing amplified copies of the selectable marker gene can be selected for by culturing the cells in the presence of the appropriate selectable agent. The activated endogenous gene is amplified in tandem with the amplifiable selectable marker gene. Cells containing many copies of the activated endogenous gene are useful for in vitro protein production and gene therapy.

Gene targeting and amplification as disclosed in the present invention are particularly useful for activating the expression of genes which form transcription units which are sufficiently large that they are difficult to

-7-

isolate and express, or for activating genes for which the entire protein coding region is unavailable or has not been cloned.

5 In a further embodiment, expression of a gene which is expressed in a cell as obtained is enhanced or caused to display a pattern of regulation or induction that is different than evident in the corresponding nontransfected cell. In another embodiment, expression of a gene which is expressed in a cell as obtained is reduced (i.e.,
10 lessened or eliminated). The present invention also describes a method by which homologous recombination is used to convert a gene into a cDNA copy, devoid of introns, for transfer into yeast or bacteria for in vitro protein production.

15 Transfected cells of the present invention are useful in a number of applications in humans and animals. In one embodiment, the cells can be implanted into a human or an animal for protein delivery in the human or animal. For example, hGH, hEPO, human insulinotropin, and other proteins can be delivered systemically or locally in humans
20 for therapeutic benefits. In addition, transfected non-human cells producing growth hormone, erythropoietin, insulinotropin and other proteins of non-human origin may be produced.

25 Barrier devices, which contain transfected cells which express a therapeutic product and through which the therapeutic product is freely permeable, can be used to retain cells in a fixed position in vivo or to protect and isolate the cells from the host's immune system. Barrier
30 devices are particularly useful and allow transfected immortalized cells, transfected xenogeneic cells, or transfected allogeneic cells to be implanted for treatment of human or animal conditions or for agricultural uses (e.g., bovine growth hormone for dairy production).
35 Barrier devices also allow convenient short-term (i.e.,

-8-

transient) therapy by providing ready access to the cells for removal when the treatment regimen is to be halted for any reason. In addition, transfected xenogeneic and allogeneic cells may be used in the absence of barrier devices for short-term gene therapy, such that the gene product produced by the cells will be delivered in vivo until the cells are rejected by the host's immune system.

Transfected cells of the present invention are also useful for eliciting antibody production or for immunizing humans and animals against pathogenic agents. Implanted transfected cells can be used to deliver immunizing antigens that result in stimulation of the host's cellular and humoral immune responses. These immune responses can be designed for protection of the host from future infectious agents (i.e., for vaccination), to stimulate and augment the disease-fighting capabilities directed against an ongoing infection, or to produce antibodies directed against the antigen produced in vivo by the transfected cells that can be useful for therapeutic or diagnostic purposes. Removable barrier devices containing the cells can be used to allow a simple means of terminating exposure to the antigen. Alternatively, the use of cells that will ultimately be rejected (xenogeneic or allogeneic transfected cells) can be used to limit exposure to the antigen, since antigen production will cease when the cells have been rejected.

The methods of the present invention can be used to produce primary, secondary, or immortalized cells producing a wide variety of therapeutically useful products, including (but not limited to): hormones, cytokines, antigens, antibodies, enzymes, clotting factors, transport proteins, receptors, regulatory proteins, structural proteins, transcription factors, ribozymes or anti-sense RNA. Additionally, the methods of the present invention can be used to produce cells which produce non-naturally

-9-

occurring ribozymes, proteins, or nucleic acids which are useful for in vitro production of a therapeutic product or for gene therapy.

Brief Description of the Drawings

5 Figure 1 is a schematic diagram of a strategy for transcriptionally activating the hEPO gene; thick lines, mouse metallothionein I promoter; stippled box, 5' untranslated region of hGH; solid box, hGH exon 1; striped box, 10 bp splice-donor sequence from hEPO intron 1; 10 cross-hatched box, 5' untranslated region of hEPO; open numbered boxes, hEPO coding sequences; diagonally-stripped box, hEPO 3' untranslated sequences; HIII, HindIII site.

Figure 2 is a schematic diagram of a strategy for transcriptionally activating the hEPO gene; thick lines, 15 mouse metallothionein I promoter; stippled box, 5' untranslated region of hGH; solid box, hGH exon 1; open numbered boxes, hEPO coding sequences; diagonally-stripped box, hEPO 3' untranslated sequences; HIII, HindIII site.

Figure 3 is a schematic representation of plasmid 20 pXGH5, which includes the hGH gene under the control of the mouse metallothionein promoter.

Figure 4 is a schematic representation of plasmid pE3neoEPO. The positions of the human erythropoietin gene and the neomycin phosphotransferase gene (neo) and 25 ampicillin (amp) resistance genes are indicated. Arrows indicate the directions of transcription of the various genes. pmMT1 denotes the mouse metallothionein promoter (driving hEPO expression) and pTK denotes the Herpes Simplex Virus thymidine kinase promoter (driving neo 30 expression). The dotted regions of the map mark the positions of sequences derived from the human hypoxanthine-guanine phosphoribosyl transferase (HPRT) locus. The relative positions of restriction endonuclease recognition sites are indicated.

-10-

Figure 5 is a schematic representation of plasmid pCDNEO, which includes the neo coding region (BamHI-BglII fragment) from plasmid pSV2neo inserted into the BamHI site of plasmid pcD; the Amp-R and pBR322Ori sequences from pBR322; and the polyA, 16S splice junctions and early promoter regions from SV40.

Figure 6 is a schematic representation of plasmid pREPO4.

Figure 7 is a graphic representation of erythropoietin expression in a targeted human cell line subjected to stepwise selection in 0.02, 0.05, 0.1, 0.2 and 0.4 μ M methotrexate.

Figure 8 is a schematic representation of plasmid pREPO15. Fragments derived from genomic hEPO sequences are indicated by filled boxes. The region between BamHI (3537) and BgIII'/HindIII' corresponds to sequences at positions 1-4008 in Genbank entry HUMERPALU. The region between BgIII'/HindIII' (11463) corresponds to DNA sequences at positions 4009-5169 in Genbank entry HUMERPALU. The region between HindIII (11463) and XhoI (624) contains sequence corresponding to positions 7-624 of Genbank entry HUMERPA. CMV promoter sequences are shown as an open box and contains sequence from nucleotides 546-2105 of Genbank sequence HS5MIEP. The neo gene is shown as an open box with an arrow. The thymidine kinase (tk) promoter driving the neo gene is shown as a hatched box. pBSIISK+ sequences including the amp gene are indicated by a thin line.

Figure 9A presents restriction enzyme maps and schematic representations of the products observed upon digestion of the endogenous hEPO gene (top) and the activated hEPO gene after homologous recombination with the targeting fragment from pREPO15 (bottom).

Figure 9B presents the results of restriction enzyme digestion and Southern hybridization analysis of untreated

-11-

(HF) and targeted (T1) human fibroblast clone HF342-15 (see Example 7).

Figure 10 is a schematic representation of plasmid pREPO18. Fragments derived from genomic hEPO sequences are indicated by filled boxes. The region between BamHI (3537) and ClaI (7554) corresponds to sequences at positions 1-4008 in Genbank entry HUMERPALU. The region between ATG (12246) and HindIII (13426) corresponds to DNA sequence at positions 4009-5169 in Genbank entry HUMERPLAU. The region between HindIII (13426) and XhoI (624) contains sequence corresponding to positions 7-624 of Genbank entry HUMERPA. CMV promoter sequences are shown as an open box and contains sequence from nucleotides 546-2015 of Genbank sequence HS5MIEP. The dihydrofolate reductase (dhfr) transcription unit is shown as a stippled box with an arrow. The neo gene is shown as an open box with an arrow. The tk promoter driving the neo gene is shown as a hatched box. pBSIISK+ sequences including the amp gene are indicated by a thin line.

Figure 11 is a schematic illustration of a construct of the invention for activating and amplifying an intronless gene, the α -interferon gene, where the construct comprises a first targeting sequence (1), an amplifiable marker gene (AM), a selectable marker gene (SM), a regulatory sequence, a CAP site, a splice-donor site (SD), an intron (thin lines), a splice-acceptor site (SA) and a second targeting sequence (2). The black box represents coding DNA and the stippled boxes represent untranslated sequences.

Figure 12 is a schematic illustration of a construct of the invention for activating and amplifying an endogenous gene wherein the first exon contributes to the signal peptide, the human GM-CSF gene, where the construct comprises a first targeting sequence (1), an amplifiable marker gene (AM), a selectable marker gene (SM), a regula-

-12-

tory sequence, a CAP site, a splice-donor site (SD), and a second targeting sequence (2). The black boxes represent coding DNA and the stippled boxes represent untranslated sequences.

- 5 Figure 13 is a schematic illustration of a construct of the invention for activating and amplifying an endogenous gene wherein the first exon contributes to the signal peptide, the human G-CSF gene, where the construct comprises a first targeting sequence (1), an amplifiable
- 10 marker gene (AM), a selectable marker gene (SM), a regulatory sequence, a CAP site, a splice-donor site (SD), and a second targeting sequence (2). The black boxes represent coding DNA and the stippled boxes represent untranslated sequences.
- 15 Figure 14 is a schematic illustration of a construct of the invention for activating and amplifying an endogenous gene wherein the first exon is non-coding, the human FSH β gene, where the construct comprises a first targeting sequence (1), an amplifiable marker gene (AM), a selecta-
- 20 ble marker gene (SM), a regulatory sequence, a CAP site, a splice-donor site (SD), and a second targeting sequence (2). The black boxes represent coding DNA and the stippled boxes represent untranslated sequences.

Detailed Description of the Invention

- 25 The invention is based upon the discovery that the regulation or activity of endogenous genes of interest in a cell can be altered by inserting into the cell genome, at a preselected site, through homologous recombination, DNA constructs comprising: (a) a targeting sequence; (b) a
- 30 regulatory sequence; (c) an exon and (d) an unpaired splice-donor site, wherein the targeting sequence directs the integration of elements (a) - (d) such that the elements (b) - (d) are operatively linked to the endogenous gene. In another embodiment, the DNA constructs comprise:

-13-

(a) a targeting sequence, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a) - (f) such that the elements of (b) - (f) are operatively linked to the first exon of the endogenous gene. The targeting sequences used are selected with reference to the site into which the DNA is to be inserted. In both embodiments the targeting event is used to create a new transcription unit, which is a fusion product of sequences introduced by the targeting DNA constructs and the endogenous cellular gene. As discussed herein, for example, the formation of the new transcription unit allows transcriptionally silent genes (genes not expressed in a cell prior to transfection) to be activated in host cells by introducing into the host cell's genome DNA constructs of the present invention. As also discussed herein, the expression of an endogenous gene which is expressed in a cell as obtained can be altered in that it is increased, reduced, including eliminated, or the pattern of regulation or induction may be changed through use of the method and DNA constructs of the present invention.

The present invention as set forth above, relates to a method of gene or DNA targeting in cells of eukaryotic origin, such as of fungal, plant or animal, such as, vertebrate, particularly mammalian, and even more particularly human origin. That is, it relates to a method of introducing DNA into a cell, such as primary, secondary, or immortalized cells of vertebrate origin, through homologous recombination or targeting of the DNA, which is introduced into genomic DNA of the cells at a preselected site. It is particularly related to homologous recombination in which the transcription and/or translation products of endogenous genes are modified through the use of DNA constructs comprising a targeting sequence, a regula-

-14-

tory sequence, an exon and a splice-donor site. The present invention further relates to homologously recombinant cells produced by the present method and to uses of the homologously recombinant cells.

5 The present invention also relates to a method of activating a gene which is present in primary cells, secondary cells or immortalized cells of vertebrate origin, but is normally not expressed in the cells. Homologous recombination or targeting is used to introduce into
10 the cell's genome sequences which causes the gene to be expressed in the recipient cell. In a further embodiment, expression of a gene in a cell is enhanced or the pattern of regulation or induction of a gene is altered, through introduction of the DNA construct. As a result, the
15 encoded product is expressed at levels higher than evident in the corresponding nontransfected cell. The present method and DNA constructs are also useful to produce cells in which expression of a desired product is less in the transfected cell than in the corresponding nontransfected
20 cell. That is, in the transfected cell, less protein (including no protein) is produced than in the cells as obtained.

In another embodiment, a normally silent gene encoding a desired product is activated in a transfected,
25 primary, secondary, or immortalized cell and amplified. This embodiment is a method of introducing, by homologous recombination with genomic DNA, DNA sequences which are not normally functionally linked to the endogenous gene and (1) which, when inserted into the host genome at or
30 near the endogenous gene, serve to alter (e.g., activate) the expression of the endogenous gene, and further (2) allow for selection of cells in which the activated endogenous gene is amplified. Alternatively, expression of a gene normally expressed in the cell as obtained is enhanced and the gene is amplified.
35

-15-

The following is a description of the DNA constructs of the present invention, methods in which they are used to produce transfected cells, transfected cells and uses of these cells.

5 The DNA Construct

The DNA construct of the present invention includes at least the following components: a targeting sequence; a regulatory sequence; an exon and an unpaired splice-donor site. In the construct, the exon is 3' of the
10 regulatory sequence and the unpaired splice-donor site is 3' of the exon. In addition, there can be multiple exons and/or introns preceding (5' to) the exon flanked by the unpaired splice-donor site. As described herein, there frequently are additional construct components, such as a
15 selectable markers or amplifiable markers.

The DNA in the construct may be referred to as exogenous. The term "exogenous" is defined herein as DNA which is introduced into a cell by the method of the present invention, such as with the DNA constructs defined herein.
20 Exogenous DNA can possess sequences identical to or different from the endogenous DNA present in the cell prior to transfection.

The Targeting Sequence or Sequences

The targeting sequence or sequences are DNA sequences
25 which permit legitimate homologous recombination into the genome of the selected cell containing the gene of interest. Targeting sequences are, generally, DNA sequences which are homologous to (i.e., identical or sufficiently similar to cellular DNA such that the targeting sequence
30 and cellular DNA can undergo homologous recombination) DNA sequences normally present in the genome of the cells as obtained (e.g., coding or noncoding DNA, lying upstream of the transcriptional start site, within, or downstream of

-16-

the transcriptional stop site of a gene of interest, or sequences present in the genome through a previous modification). The targeting sequence or sequences used are selected with reference to the site into which the DNA in
5 the DNA construct is to be inserted.

One or more targeting sequences can be employed. For example, a circular plasmid or DNA fragment preferably employs a single targeting sequence. A linear plasmid or DNA fragment preferably employs two targeting sequences.
10 The targeting sequence or sequences can, independently, be within the gene of interest (such as, the sequences of an exon and/or intron), immediately adjacent to the gene of interest (i.e., with no additional nucleotides between the targeting sequence and the coding region of the gene of
15 interest), upstream gene of interest (such as the sequences of the upstream non-coding region or endogenous promoter sequences), or upstream of and at a distance from the gene (such as, sequences upstream of the endogenous promoter). The targeting sequence or sequences can include
20 those regions of the targeted gene presently known or sequenced and/or regions further upstream which are structurally uncharacterized but can be mapped using restriction enzymes and determined by one skilled in the art.

As taught herein, gene targeting can be used to
25 insert a regulatory sequence isolated from a different gene, assembled from components isolated from difference cellular and/or viral sources, or synthesized as a novel regulatory sequence by genetic engineering methods within, immediately adjacent to, upstream, or at a substantial
30 distance from an endogenous cellular gene. Alternatively or additionally, sequences which affect the structure or stability of the RNA or protein produced can be replaced, removed, added, or otherwise modified by targeting. For example, RNA stability elements, splice sites, and/or
35 leader sequences of RNA molecules can be modified to

-17-

improve or alter the function, stability, and/or translatability of an RNA molecule. Protein sequences may also be altered, such as signal sequences, propeptide sequences, active sites, and/or structural sequences for enhancing or
5 modifying transport, secretion, or functional properties of a protein. According to this method, introduction of the exogenous DNA results in the alteration of the normal expression properties of a gene and/or the structural properties of a protein or RNA.

10 The Regulatory Sequence

The regulatory sequence of the DNA construct can be comprised of one or more promoters (such as a constitutive or inducible promoter), enhancers, scaffold-attachment regions or matrix attachment sites, negative regulatory
15 elements, transcription factor binding sites, or combinations of said sequences.

The regulatory sequence can contain an inducible promoter, with the result that cells as produced or as introduced into an individual do not express the product
20 but can be induced to do so (i.e., expression is induced after the transfected cells are produced but before implantation or after implantation). DNA encoding the desired product can, of course, be introduced into cells in such a manner that it is expressed upon introduction
25 (e.g., under a constitutive promoter). The regulatory sequence can be isolated from cellular or viral genomes, (such regulatory sequences include those that regulate the expression of SV40 early or late genes, adenovirus major late genes, the mouse metallothionein-I gene, the elongation factor-1 α gene, cytomegalovirus genes, collagen
30 genes, actin genes, immunoglobulin genes or the HMG-CoA reductase gene). The regulatory sequence preferably contains transcription factor binding sites, such as a TATA Box, CCAAT Box, AP1, Sp1 or NF- κ B binding sites.

-18-

Additional DNA Construct Elements

The DNA construct further comprises one or more exons. An exon is defined herein as a DNA sequence which is copied into RNA and is present in a mature mRNA molecule. The exons can, optionally, contain DNA which encodes one or more amino acids and/or partially encodes an amino acid (i.e., one or two bases of a codon). Alternatively, the exon contains DNA which corresponds to a 5' non-coding region. Where the exogenous exon or exons encode one or more amino acids and/or a portion of an amino acid, the DNA construct is designed such that, upon transcription and splicing, the reading frame is in-frame with the second exon or coding region of the targeted gene. As used herein, in-frame means that the encoding sequences of a first exon and a second exon, when fused, join together nucleotides in a manner that does not change the appropriate reading frame of the portion of the mRNA derived from the second exon.

Where the first exon of the targeted gene contains the sequence ATG to initiate translation, the exogenous exon of the construct preferably contains an ATG and, if required, one or more nucleotides such that the resulting coding region of the mRNA including the second and subsequent exons of the targeted gene is in-frame. Examples of such targeted genes in which the first exon contains an ATG include the genes encoding hEPO, hGH, human colony stimulating factor-granulocyte/macrophage (hGM-CSF), and human colony stimulating factor-granulocyte (hG-CSF).

A splice-donor site is a sequence which directs the splicing of one exon to another exon. Typically, the first exon lies 5' of the second exon, and the splice-donor site overlapping and flanking the first exon on its 3' side recognizes a splice-acceptor site flanking the second exon on the 5' side of the second exon. Splice-donor sites have a characteristic consensus sequence

-19-

represented as: (A/C)AG GURAGU (where R denotes a purine nucleotide) with the GU in the fourth and fifth positions, being required (Jackson, I.J., *Nucleic Acids Research* 19: 3715-3798 (1991)). The first three bases of the splice-donor consensus site are the last three bases of the exon. Splice-donor sites are functionally defined by their ability to effect the appropriate reaction within the mRNA splicing pathway.

An unpaired splice-donor site is defined herein as a splice-donor site which is present in a targeting construct and is not accompanied in the construct by a splice-acceptor site positioned 3' to the unpaired splice-donor site. The unpaired splice-donor site results in splicing to an endogenous splice-acceptor site.

A splice-acceptor site in a sequence which, like a splice-donor site, directs the splicing of one exon to another exon. Acting in conjunction with a splice-donor site, the splicing apparatus uses a splice-acceptor site to effect the removal of an intron. Splice-acceptor sites have a characteristic sequence represented as: YYYYYYYY-NYAG, where Y denotes any pyrimidine and N denotes any nucleotide (Jackson, I.J., *Nucleic Acids Research* 19:3715-3798 (1991)).

An intron is defined as a sequence of one or more nucleotides lying between two exons and which is removed, by splicing, from a precursor RNA molecule in the formation of an mRNA molecule.

The regulatory sequence is, for example, operatively linked to an ATG start codon, which initiates translation. Optionally, a CAP site (a specific mRNA initiation site which is associated with and utilized by the regulatory region) is operatively linked to the regulatory sequence and the ATG start codon. Alternatively, the CAP site associated with and utilized by the regulatory sequence is not included in the targeting construct, and the trans-

-20-

criptional apparatus will define a new CAP site. For most genes, a CAP site is usually found approximately 25 nucleotides 3' of the TATA box. In one embodiment, the splice-donor site is placed immediately adjacent to the ATG, for example, where the presence of one or more nucleotides is not required for the exogenous exon to be in-frame with the second exon of the targeted gene. Preferably, DNA encoding one or more amino acids or portions of an amino acid in-frame with the coding sequence of the targeted gene, is placed immediately adjacent to the ATG on its 3' side. In such an embodiment, the splice-donor site is placed immediately adjacent to the encoding DNA on its 3' side.

Operatively linked or functionally placed is defined as a configuration in which the exogenous regulatory sequence, exon, splice-donor site and, optionally, a sequence and splice-acceptor site are appropriately targeted at a position relative to an endogenous gene such that the regulatory element directs the production of a primary RNA transcript which initiates at a CAP site (optionally included in the targeting construct) and includes sequences corresponding to the exon and splice-donor site of the targeting construct, DNA lying upstream of the endogenous gene's regulatory region (if present), the endogenous gene's regulatory region (if present), the endogenous genes 5' nontranscribed region (if present), and exons and introns (if present) of the endogenous gene. In an operatively linked configuration the splice-donor site of the targeting construct directs a splicing event to a splice-acceptor site flanking one of the exons of the endogenous gene, such that a desired protein can be produced from the fully spliced mature transcript. In one embodiment, the splice-acceptor site is endogenous, such that the splicing event is directed to an endogenous exon, for example, of the endogenous gene. In another embodiment where the

-21-

splice-acceptor site is included in the targeting construct, the splicing event removes the intron introduced by the targeting construct.

The encoding DNA (e.g., in exon 1 of the targeting
5 construct) employed can optionally encode one or more
amino acids, and/or a portion of an amino acid, which are
the same as those of the endogenous protein. The encoding
DNA sequence employed herein can, for example, correspond
10 to the first exon of the gene of interest. The
encoding DNA can alternatively encode one or more amino
acids or a portion of an amino acid different from the
first exon of the protein of interest. Such an embodiment
is of particular interest where the amino acids of the
15 first exon of the protein of interest are not critical to
the activity or activities of the protein. For example,
when fusions to the endogenous hEPO gene are constructed,
sequences encoding the first exon of hGH can be employed.
In this example, fusion of hGH exon 1 to hEPO exon 2
20 results in the formation of a hybrid signal peptide which
is functional. In related constructs, any exon of human
or non-human origin in which the encoded amino acids do
not prevent the function of the hybrid signal peptide can
be used. In a related embodiment, this technique can also
be employed to correct a mutation found in a target gene.

25 Where the desired product is a fusion protein of the
endogenous protein and encoding sequences in the targeting
construct, the exogenous encoding DNA incorporated into
the cells by the present method includes DNA which encodes
one or more exons or a sequence of cDNA corresponding to a
30 translation or transcription product which is to be fused
to the product of the endogenous targeted gene. In this
embodiment, targeting is used to prepare chimeric or
multifunctional proteins which combine structural, enzymatic,
or ligand or receptor binding properties from two
35 or more proteins into one polypeptide. For example, the

-22-

exogenous DNA can encode an anchor to the membrane for the targeted protein or a signal peptide to provide or improve cellular secretion, leader sequences, enzymatic regions, transmembrane domain regions, co-factor binding regions or other functional regions. Examples of proteins which are not normally secreted, but which could be fused to a signal protein to provide secretion include dopa-decarboxylase, transcriptional regulatory proteins, α -galactosidase and tyrosine hydroxylase.

Where the first exon of the targeted gene corresponds to a non-coding region (for example, the first exon of the follicle-stimulating hormone beta (FSH β) gene, an exogenous ATG is not required and, preferably, is omitted.

The DNA of the construct can be obtained from sources in which it occurs in nature or can be produced, using genetic engineering techniques or synthetic processes.

The Targeted Gene and Resulting Product

The DNA construct, when transfected into cells, such as primary, secondary or immortalized cells, can control the expression of a desired product for example, the active or, functional portion of the protein or RNA. The product can be, for example, a hormone, a cytokine, an antigen, an antibody, an enzyme, a clotting factor, a transport protein, a receptor, a regulatory protein, a structural protein, a transcription factor, an anti-sense RNA, or a ribozyme. Additionally, the product can be a protein or a nucleic acid which does not occur in nature (i.e., a fusion protein or nucleic acid).

The method as described herein can produce one or more therapeutic products, such as erythropoietin, calcitonin, growth hormone, insulin, insulinotropin, insulin-like growth factors, parathyroid hormone, interferon β , and interferon β , nerve growth factors, FSH β , TGF- β , tumor necrosis factor, glucagon, bone growth factor-2, bone

-23-

growth factor-7, TSH- β , interleukin 1, interleukin 2, interleukin 3, interleukin 6, interleukin 11, interleukin 12, CSF-granulocyte, CSF-macrophage, CSF-granulocyte/macrophage, immunoglobulins, catalytic antibodies, protein kinase C, glucocerebrosidase, superoxide dismutase, tissue plasminogen activator, urokinase, antithrombin III, DNase, α -galactosidase, tyrosine hydroxylase, blood clotting factors V, blood clotting factor VII, blood clotting factor VIII, blood clotting factor IX, blood clotting factor X, blood clotting factor XIII, apolipoprotein E or apolipoprotein A-I, globins, low density lipoprotein receptor, IL-2 receptor, IL-2 antagonists, alpha-1 anti-trypsin, immune response modifiers, and soluble CD4.

Selectable Markers and Amplification

The identification of the targeting event can be facilitated by the use of one or more selectable marker genes. These markers can be included in the targeting construct or be present on different constructs. Selectable markers can be divided into two categories: positively selectable and negatively selectable (in other words, markers for either positive selection or negative selection). In positive selection, cells expressing the positively selectable marker are capable of surviving treatment with a selective agent (such as neo, xanthine-guanine phosphoribosyl transferase (gpt), dhfr, adenosine deaminase (ada), puromycin (pac), hygromycin (hyg), CAD which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydro-orotase glutamine synthetase (GS), multidrug resistance 1 (mdr1) and histidine D (hisD), allowing for the selection of cells in which the targeting construct integrated into the host cell genome. In negative selection, cells expressing the negatively selectable marker are destroyed in the presence of the selective agent. The identification of the targeting

-24-

event can be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker (Mansour, S.L. *et al.*, *Nature* 336:348-352 (1988)). Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial gpt gene.

A variety of selectable markers can be incorporated into primary, secondary or immortalized cells. For example, a selectable marker which confers a selectable phenotype such as drug resistance, nutritional auxotrophy, resistance to a cytotoxic agent or expression of a surface protein, can be used. Selectable marker genes which can be used include neo, gpt, dhfr, ada, pac, hyg, CAD, GS, mdrl and hisD. The selectable phenotype conferred makes it possible to identify and isolate recipient cells.

Amplifiable genes encoding selectable markers (e.g., ada, GS, dhfr and the multifunctional CAD gene) have the added characteristic that they enable the selection of cells containing amplified copies of the selectable marker inserted into the genome. This feature provides a mechanism for significantly increasing the copy number of an adjacent or linked gene for which amplification is desirable. Mutated versions of these sequences showing improved selection properties and other amplifiable sequences can also be used.

The order of components in the DNA construct can vary. Where the construct is a circular plasmid, the order of elements in the resulting structure can be: targeting sequence - plasmid DNA (comprised of sequences

-25-

used for the selection and/or replication of the targeting plasmid in a microbial or other suitable host) - selectable marker(s) - regulatory sequence - exon - splice-donor site. Preferably, the plasmid containing the targeting

5 sequence and exogenous DNA elements is cleaved with a restriction enzyme that cuts one or more times within the targeting sequence to create a linear or gapped molecule prior to introduction into a recipient cell, such that the free DNA ends increase the frequency of the desired homologous recombination event as described herein. In addition,

10 the free DNA ends may be treated with an exonuclease to create protruding 5' or 3' overhanging single-stranded DNA ends to increase the frequency of the desired homologous recombination event. In this embodiment, homologous

15 recombination between the targeting sequence and the cellular target will result in two copies of the targeting sequences, flanking the elements contained within the introduced plasmid.

Where the construct is linear, the order can be, for example: a first targeting sequence - selectable marker - regulatory sequence - an exon - a splice-donor site - a second targeting sequence or, in the alternative, a first targeting sequence - regulatory sequence - an exon - a splice-donor site - DNA encoding a selectable marker - a

25 second targeting sequence. Cells that stably integrate the construct will survive treatment with the selective agent; a subset of the stably transfected cells will be homologously recombinant cells. The homologously recombinant cells can be identified by a variety of techniques,

30 including PCR, Southern hybridization and phenotypic screening.

In another embodiment, the order of the construct can be: a first targeting sequence - selectable marker - regulatory sequence - an exon - a splice-donor site - an

35 intron - a splice-acceptor site - a second targeting sequence.

-26-

Alternatively, the order of components in the DNA construct can be, for example: a first targeting sequence - selectable marker 1 - regulatory sequence - an exon - a splice-donor site - a second targeting sequence - selectable marker 2, or, alternatively, a first targeting sequence - regulatory sequence - an exon - a splice-donor site - selectable marker 1 - a second targeting sequence - selectable marker 2. In this embodiment selectable marker 2 displays the property of negative selection. That is, the gene product of selectable marker 2 can be selected against by growth in an appropriate media formulation containing an agent (typically a drug or metabolite analog) which kills cells expressing selectable marker 2. Recombination between the targeting sequences flanking selectable marker 1 with homologous sequences in the host cell genome results in the targeted integration of selectable marker 1, while selectable marker 2 is not integrated. Such recombination events generate cells which are stably transfected with selectable marker 1 but not stably transfected with selectable marker 2, and such cells can be selected for by growth in the media containing the selective agent which selects for selectable marker 1 and the selective agent which selects against selectable marker 2.

The DNA construct also can include a positively selectable marker that allows for the selection of cells containing amplified copies of that marker. The amplification of such a marker results in the co-amplification of flanking DNA sequences. In this embodiment, the order of construct components is, for example: a first targeting sequence - an amplifiable positively selectable marker - a second selectable marker (optional) - regulatory sequence - an exon - a splice-donor site - a second targeting DNA sequence.

-27-

In this embodiment, the activated gene can be further amplified by the inclusion of a selectable marker gene which has the property that cells containing amplified copies of the selectable marker gene can be selected for by culturing the cells in the presence of the appropriate selectable agent. The activated endogenous gene will be amplified in tandem with the amplified selectable marker gene. Cells containing many copies of the activated endogenous gene may produce very high levels of the desired protein and are useful for in vitro protein production and gene therapy.

In any embodiment, the selectable and amplifiable marker genes do not have to lie immediately adjacent to each other.

Optionally, the DNA construct can include a bacterial origin of replication and bacterial antibiotic resistance markers or other selectable markers, which allow for large-scale plasmid propagation in bacteria or any other suitable cloning/host system. A DNA construct which includes DNA encoding a selectable marker, along with additional sequences, such as a promoter, and splice junctions, can be used to confer a selectable phenotype upon transfected cells (e.g., plasmid pcDNEO, schematically represented in Figure 4). Such a DNA construct can be co-transfected into primary or secondary cells, along with a targeting DNA sequence, using methods described herein.

Transfection and Homologous Recombination

According to the present method, the construct is introduced into the cell, such as a primary, secondary, or immortalized cell, as a single DNA construct, or as separate DNA sequences which become incorporated into the chromosomal or nuclear DNA of a transfected cell.

The targeting DNA construct, including the targeting sequences, regulatory sequence, an exon, a splice-donor

-28-

site and selectable marker gene(s), can be introduced into cells on a single DNA construct or on separate constructs. The total length of the DNA construct will vary according to the number of components (targeting sequences, regulatory sequences, exons, selectable marker gene, and other elements, for example) and the length of each. The entire construct length will generally be at least about 200 nucleotides. Further, the DNA can be introduced as linear, double-stranded (with or without single-stranded regions at one or both ends), single-stranded, or circular.

Any of the construct types of the disclosed invention is then introduced into the cell to obtain a transfected cell. The transfected cell is maintained under conditions which permit homologous recombination, as is known in the art (Capecchi, M.R., Science 244:1288-1292 (1989)). When the homologously recombinant cell is maintained under conditions sufficient for transcription of the DNA, the regulatory region introduced by the targeting construct, as in the case of a promoter, will activate transcription.

The DNA constructs may be introduced into cells by a variety of physical or chemical methods, including electroporation, microinjection, microprojectile bombardment, calcium phosphate precipitation, and liposome-, polybrene-, or DEAE dextran-mediated transfection. Alternatively, infectious vectors, such as retroviral, herpes, adenovirus, adenovirus-associated, mumps and poliovirus vectors, can be used to introduce the DNA.

Optionally, the targeting DNA can be introduced into a cell in two or more separate DNA fragments. In the event two fragments are used, the two fragments share DNA sequence homology (overlap) at the 3' end of one fragment and the 5' end of the other, while one carries a first targeting sequence and the other carries a second targeting sequence. Upon introduction into a cell, the two

-29-

fragments can undergo homologous recombination to form a single fragment with the first and second targeting sequences flanking the region of overlap between the two original fragments. The product fragment is then in a form suitable for homologous recombination with the cellular target sequences. More than two fragments can be used, designed such that they will undergo homologous recombination with each other to ultimately form a product suitable for homologous recombination with the cellular target sequences as described above.

The Homologously Recombinant Cells

The targeting event results in the insertion of the regulatory sequence of the targeting construct, placing the endogenous gene under their control (for example, by insertion of either a promoter or an enhancer, or both, upstream of the endogenous gene or regulatory region). Optionally, the targeting event can simultaneously result in the deletion of the endogenous regulatory element, such as the deletion of a tissue-specific negative regulatory element. The targeting event can replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally-occurring elements, or displays a pattern of regulation or induction that is different from the corresponding nontransfected cell. In this embodiment the naturally occurring sequences are deleted and new sequences are added. Alternatively, the endogenous regulatory elements are not removed or replaced but are disrupted or disabled by the targeting event, such as by targeting the exogenous sequences within the endogenous regulatory elements.

After the DNA is introduced into the cell, the cell is maintained under conditions appropriate for homologous recombination to occur between the genomic DNA and a

-30-

portion of the introduced DNA, as is known in the art (Capecchi, M.R., Science 244:1288-1292 (1989)).

Homologous recombination between the genomic DNA and the introduced DNA results in a homologously recombinant
5 cell, such as a fungal, plant or animal, and particularly, primary, secondary, or immortalized human or other mammalian cell in which sequences which alter the expression of an endogenous gene are operatively linked to an endogenous gene encoding a product, producing a new transcription
10 unit with expression and/or coding potential that is different from that of the endogenous gene. Particularly, the invention includes a homologously recombinant cell comprising regulatory sequences and an exon, flanked by a splice-donor site, which are introduced at a predetermined
15 site by a targeting DNA construct, and are operatively linked to the second exon of an endogenous gene. Optionally, there may be multiple exogenous exons (coding or non-coding) and introns operatively linked to any exon of the endogenous gene. The resulting homologously recombi-
20 nant cells are cultured under conditions which select for amplification, if appropriate, of the DNA encoding the amplifiable marker and the novel transcriptional unit. With or without amplification, cells produced by this method can be cultured under conditions, as are known in
25 the art, suitable for the expression of the protein, thereby producing the protein in vitro, or the cells can be used for in vivo delivery of a therapeutic protein (i.e., gene therapy).

As used herein, the term primary cell includes cells
30 present in a suspension of cells isolated from a vertebrate tissue source (prior to their being plated, i.e., attached to a tissue culture substrate such as a dish or flask), cells present in an explant derived from tissue, both of the previous types of cells plated for the first
35 time, and cell suspensions derived from these plated

-31-

cells. The term secondary cell or cell strain refers to cells at all subsequent steps in culturing. That is, the first time a plated primary cell is removed from the culture substrate and replated (passaged), it is referred to herein as a secondary cell, as are all cells in subsequent passages. Secondary cells are cell strains which consist of secondary cells which have been passaged one or more times. A cell strain consists of secondary cells that: 1) have been passaged one or more times; 2) exhibit a finite number of mean population doublings in culture; 3) exhibit the properties of contact-inhibited, anchorage dependent growth (anchorage-dependence does not apply to cells that are propagated in suspension culture); and 4) are not immortalized.

Immortalized cells are cell lines (as opposed to cell strains with the designation "strain" reserved for primary and secondary cells), a critical feature of which is that they exhibit an apparently unlimited lifespan in culture.

Cells selected for the subject method can fall into four types or categories: 1) cells which do not, as obtained, make or contain the protein or product (such as a protein that is not normally expressed by the cell or a fusion protein not normally found in nature), 2) cells which make or contain the protein or product but in quantities other than that desired (such as, in quantities less than the physiologically normal lower level for the cell as it is obtained), 3) cells which make the protein or product at physiologically normal levels for the cell as it is obtained, but are to be augmented or enhanced in their content or production, and 4) cells in which it is desirable to change the pattern of regulation or induction of a gene encoding a protein.

Primary, secondary and immortalized cells to be transfected by the present method can be obtained from a variety of tissues and include all cell types which can be

-32-

maintained in culture. For example, primary and secondary cells which can be transfected by the present method include fibroblasts, keratinocytes, epithelial cells (e.g., mammary epithelial cells, intestinal epithelial cells), endothelial cells, glial cells, neural cells, formed elements of the blood (e.g., lymphocytes, bone marrow cells), muscle cells and precursors of these somatic cell types. Where the homologously recombinant cells are to be used in gene therapy, primary cells are preferably obtained from the individual to whom the transfected primary or secondary cells are administered. However, primary cells can be obtained from a donor (other than the recipient) of the same species.

Homologously recombinant immortalized cells can also be produced by the present method and used for either protein production or gene therapy. Examples of immortalized human cell lines useful for protein production or gene therapy by the present method include, but are not limited to, HT1080 cells (ATCC CCL 121), HeLa cells and derivatives of HeLa cells (ATCC CCL 2, 2.1 and 2.2), MCF-7 breast cancer cells (ATCC BTH 22), K-562 leukemia cells (ATCC CCL 243), KB carcinoma cells (ATCC CCL 17), 2780AD ovarian carcinoma cells (Van der Blick, A.M. et al., Cancer Res. 48:5927-5932 (1988), Raji cells (ATCC CCL 86), Jurkat cells (ATCC TIB 152), Namalwa cells (ATCC CRL 1432), HL-60 cells (ATCC CCL 240), Daudi cells (ATCC CCL 213), RPMI 8226 cells (ATCC CCL 155), U-937 cells (ATCC CRL 1593), Bowes Melanoma cells (ATCC CRL 9607), WI-38VA13 subline 2R4 cells (ATCC CLL 75.1), and MOLT-4 cells (ATCC CRL 1582), as well as heterohybridoma cells produced by fusion of human cells and cells of another species.

Secondary human fibroblast strains, such as WI-38 (ATCC CCL 75) and MRC-5 (ATCC CCL 171) may be used. In addition, primary, secondary, or immortalized human cells, as well as primary, secondary, or immortalized cells from

-33-

other species which display the properties of gene amplification in vitro can be used for in vitro protein production or gene therapy.

Method of Converting a Gene into a cDNA Copy

5 The present invention also relates to a method by which homologous recombination is used to convert a gene into a cDNA copy (a gene copy devoid of introns). The cDNA copy can be transferred into yeast or bacteria for in vitro protein production, or the cDNA copy can be inserted
10 into a mammalian cell for in vitro or in vivo protein production. If the cDNA is to be transferred to microbial cells, two DNA constructs containing targeting sequences are introduced by homologous recombination, one construct upstream of and one construct downstream of a human gene
15 encoding a therapeutic protein. For example, the sequences introduced upstream include DNA sequences homologous to genomic DNA sequences at or upstream of the DNA encoding the first amino acid of a mature, processed therapeutic protein; a retroviral long term repeat (LTR); sequences
20 encoding a marker for selection in microbial cells; a regulatory element that functions in microbial cells; and DNA encoding a leader peptide that promotes secretion from microbial cells with a splice-donor site. The sequences introduced upstream are introduced near to and upstream of
25 genomic DNA encoding the first amino acid of a mature, processed therapeutic protein. The sequences introduced downstream include DNA sequences homologous to genomic DNA sequences at or downstream of the DNA encoding the last amino acid of a mature, processed protein; a microbial
30 transcriptional termination sequence; sequences capable of directing DNA replication in microbial cells; and a retroviral LTR. The sequences introduced downstream are introduced adjacent to and downstream of the DNA encoding the stop codon of the mature, processed therapeutic protein.

-34-

After the two DNA constructs are introduced into cells, the resulting cells are maintained under conditions appropriate for homologous recombination between the introduced DNA and genomic DNA, thereby producing homologously recombinant cells. Optionally, one or both of the DNA constructs can encode one or more markers for either positive or negative selection of cells containing the DNA construct, and a selection step can be added to the method after one or both of the DNA constructs have been introduced into the cells. Alternatively, the sequences encoding the marker for selection in microbial cells and the sequences capable of directing DNA replication in microbial cells can both be present in either the upstream or the downstream targeting construct, or the marker for selection in microbial cells can be present in the downstream targeting construct and the sequences capable of directing DNA replication in microbial cells can be present in the upstream targeting construct. The homologously recombinant cells are then cultured under conditions appropriate for LTR directed transcription, processing and reverse transcription of the RNA product of the gene encoding the therapeutic protein. The product of reverse transcription is a DNA construct comprising an intronless DNA copy encoding the therapeutic protein, operatively linked to DNA sequences comprising the two exogenous DNA constructs described above. The intronless DNA construct produced by the present method is then introduced into a microbial cell. The microbial cell is then cultured under conditions appropriate for expression and secretion of the therapeutic protein.

In Vivo Protein Production

Homologously recombinant cells of the present invention are useful, as populations of homologously recombinant cell lines, as populations of homologously recombi-

-35-

nant primary or secondary cells, homologously recombinant clonal cell strains or lines, homologously recombinant heterogenous cell strains or lines, and as cell mixtures in which at least one representative cell of one of the

5 four preceding categories of homologously recombinant cells is present. Such cells may be used in a delivery system for treating an individual with an abnormal or undesirable condition which responds to delivery of a therapeutic product, which is either: 1) a therapeutic

10 protein (e.g., a protein which is absent, underproduced relative to the individual's physiologic needs, defective or inefficiently or inappropriately utilized in the individual; a protein with novel functions, such as enzymatic or transport functions) or 2) a therapeutic nucleic acid

15 (e.g., RNA which inhibits gene expression or has intrinsic enzymatic activity). In the method of the present invention of providing a therapeutic protein or nucleic acid, homologously recombinant primary cells, clonal cell strains or heterogenous cell strains are administered to

20 an individual in whom the abnormal or undesirable condition is to be treated or prevented, in sufficient quantity and by an appropriate route, to express or make available the protein or exogenous DNA at physiologically relevant levels. A physiologically relevant level is one which

25 either approximates the level at which the product is normally produced in the body or results in improvement of the abnormal or undesirable condition. According to an embodiment of the invention described herein, the homologously recombinant immortalized cell lines to be admin-

30 istered can be enclosed in one or more semipermeable barrier devices. The permeability properties of the device are such that the cells are prevented from leaving the device upon implantation into an animal, but the therapeutic product is freely permeable and can leave the

35 barrier device and enter the local space surrounding the

-36-

implant or enter the systemic circulation. For example, hGH, hEPO, human insulinotropin, hGM-CSF, hG-CSF, human α -interferon, or human FSH β can be delivered systemically in humans for therapeutic benefits.

5 Barrier devices are particularly useful and allow homologously recombinant immortalized cells, homologously recombinant cells from another species (homologously recombinant xenogeneic cells), or cells from a nonhisto-
10 compatibility-matched donor (homologously recombinant allogeneic cells) to be implanted for treatment of human or animal conditions or for agricultural uses (i.e., meat and dairy production). Barrier devices also allow convenient short-term (i.e., transient) therapy by providing ready access to the cells for removal when the treatment
15 regimen is to be halted for any reason.

A number of synthetic, semisynthetic, or natural filtration membranes can be used for this purpose, including, but not limited to, cellulose, cellulose acetate, nitrocellulose, polysulfone, polyvinylidene difluoride,
20 polyvinyl chloride polymers and polymers of polyvinyl chloride derivatives. Barrier devices can be utilized to allow primary, secondary, or immortalized cells from another species to be used for gene therapy in humans.

In Vitro Protein Production

25 Homologously recombinant cells from human or non-human species according to this invention can also be used for in vitro protein production. The cells are maintained under conditions, as are known in the art, which result in expression of the protein. Proteins expressed using the
30 methods described may be purified from cell lysates or cell supernatants in order to purify the desired protein. Proteins made according to this method include therapeutic proteins which can be delivered to a human or non-human animal by conventional pharmaceutical routes as is known

-37-

in the art (e.g., oral, intravenous, intramuscular, intranasal or subcutaneous). Such proteins include hGH, hEPO, and human insulinotropin, hGM-CSF, hG-CSF, FSH β or α -interferon. These cells can be immortalized, primary, or
5 secondary cells. The use of cells from other species may be desirable in cases where the non-human cells are advantageous for protein production purposes where the non-human protein is therapeutically or commercially useful, for example, the use of cells derived from salmon for the
10 production of salmon calcitonin, the use of cells derived from pigs for the production of porcine insulin, and the use of bovine cells for the production of bovine growth hormone.

Advantages

15 The methodologies, DNA constructs, cells, and resulting proteins of the invention herein possess versatility and many other advantages over processes currently employed within the art in gene targeting. The ability to activate an endogenous gene by positioning an exogenous
20 regulatory sequence at various positions ranging from immediately adjacent to the gene of interest (directly fused to the normal gene's transcribed region) to 30 kilobase pairs or further upstream of the transcribed region of an endogenous gene, or within an intron of an
25 endogenous gene, is advantageous for gene expression in cells. For example, it can be employed to position the regulatory element upstream or downstream of regions that normally silence or negatively regulate a gene. The
30 positioning of a regulatory element upstream or downstream of such a region can override such dominant negative effects that normally inhibit transcription. In addition, regions of DNA that normally inhibit transcription or have an otherwise detrimental effect on the expression of a

-38-

gene may be deleted using the targeting constructs, described herein.

Additionally, since promoter function is known to depend strongly on the local environment, a wide range of positions may be explored in order to find those local environments optimal for function. However, since, ATG start codons are found frequently within mammalian DNA (approximately one occurrence per 48 base pairs), transcription cannot simply initiate at any position upstream of a gene and produce a transcript containing a long leader sequence preceding the correct ATG start codon, since the frequent occurrence of ATG codons in such a leader sequence will prevent translation of the correct gene product and render the message useless. Thus, the incorporation of an exogenous exon, a splice-donor site, and, optionally, an intron and a splice-acceptor site into targeting constructs comprising a regulatory region allows gene expression to be optimized by identifying the optimal site for regulatory region function, without the limitation imposed by needing to avoid inappropriate ATG start codons in the mRNA produced. This provides significantly increased flexibility in the placement of the construct and makes it possible to activate a wider range of genes. The DNA constructs of the present invention are also useful, for example, in processes for making fusion proteins encoded by recombinant, or exogenous, sequences and endogenous sequences.

Gene targeting and amplification as disclosed above are particularly useful for altering on the expression of genes which form transcription units which are sufficiently large that they are difficult to isolate and express, or for turning on genes for which the entire protein coding region is unavailable or has not been cloned. Thus, the DNA constructs described above are useful for operatively linking exogenous regulatory elements to

-39-

endogenous genes in a way that precisely defines the transcriptional unit, provides flexibility in the relative positioning of exogenous regulatory elements and endogenous genes ultimately, enables a highly controlled system for obtaining and regulating expression of genes of therapeutic interest.

Explanation of the Examples

As described herein, Applicants have demonstrated that DNA can be introduced into cells, such as primary, secondary or immortalized vertebrate cells and integrated into the genome of the transfected cells by homologous recombination. They have further demonstrated that the exogenous DNA has the desired function in the homologously recombinant (HR) cells and that correctly targeted cells can be identified on the basis of a detectable phenotype conferred by the properly targeted DNA.

Applicants describe construction of a plasmid useful for targeting to a particular locus (the HPRT locus) in the human genome and selection based upon a drug resistant phenotype (Example 1a). This plasmid is designated pE3Neo and its integration into the cellular genome at the HPRT locus produces cells which have an $hprt^-$, 6-TG resistant phenotype and are also G418 resistant. As described, they have shown that pE3Neo functions properly in gene targeting in an established human fibroblast cell line (Example 1b), by demonstrating localization of the DNA introduced into established cells within exon 3 of the HPRT gene.

In addition, Applicants demonstrate gene targeting in primary and secondary human skin fibroblasts using pE3Neo (Example 1c). The subject application further demonstrates that modification of DNA termini enhances targeting of DNA into genomic DNA (Examples 1c and 1e). Applicants also describe methods by which a gene can be inserted at a preselected site in the genome of a cell,

-40-

such as a primary, secondary, or immortalized cell by gene targeting (Example 1d).

In addition, the present invention relates to a method of protein production using transfected cells. The method involves transfecting cells, such as primary cells, secondary cells or immortalized cells, with exogenous DNA which encodes a therapeutic product or with DNA which is sufficient to target to an endogenous gene which encodes a therapeutic product. For example, Examples 1g, 1h, 1j, 1k, 2, 3, 4 and 6 - 9 describe protein production by targeting of a selected endogenous gene with DNA sequence elements which will alter the expression of the endogenous gene.

Applicants also describe DNA constructs and methods for amplifying an endogenous cellular gene that has been activated by gene targeting (Examples 3, 6, 8 and 9).

Examples 1f-1h, 2, 4 and 6 illustrate embodiments in which the normal regulatory sequences upstream of the human EPO gene are altered to allow expression of hEPO in primary or secondary fibroblast strains which do not express EPO in detectable quantities in their untransfected state. In one embodiment the product of targeting leaves the normal EPO protein intact, but under the control of the mouse metallothionein promoter. Examples 1i and 1j demonstrate the use of similar targeting constructs to activate the endogenous growth hormone gene in primary or secondary human fibroblasts. In other embodiments described for activating EPO expression in human fibroblasts, the products of targeting events are chimeric transcription units, in which the first exon of the human growth hormone gene is positioned upstream of EPO exons 2-5. The product of transcription (controlled by the mouse metallothionein promoter), splicing, and translation is a protein in which amino acids 1-4 of the hEPO signal peptide are replaced with amino acid residues 1-3 of hGH.

-41-

The chimeric portion of this protein, the signal peptide, is removed prior to secretion from cells. Example 5 describes targeting constructs and methods for producing cells which will convert a gene (with introns) into an expressible cDNA copy of that gene (without introns) and the recovery of such expressible cDNA molecules in microbial (e.g., yeast or bacterial) cells. Example 6 describes construction of a targeting vector, designated pREPO4 for dual selection and selection of cells in which the *dhfr* gene is amplified. Plasmid pREPO4 has been used to amplify the human EPO (hEPO) locus in HT1080 cells (an immortalized human cell line) after activation of the endogenous hEPO gene by homologous recombination. As described, stepwise selection in methotrexate-containing media resulted in a 70-fold increase in hEPO production in cells resistant to 0.4 μ M methotrexate.

Examples 7 and 8 describe methods for inserting a regulatory sequence upstream of the normal EPO promoter and methods for EPO production using such a construct. In addition, Example 8 describes the amplification of a targeted EPO gene produced by the method of Example 7. Example 9 describes methods for targeting the human α -interferon, GM-CSF, G-CSF, and FSH β genes to create cells useful for in protein production.

The Examples provide methods for activating or for activating and amplifying endogenous genes by gene targeting which do not require manipulation or other uses of the target genes' protein coding regions. Using the methods and DNA constructs or plasmids taught herein or modifications thereof which are apparent to one of ordinary skill in the art, gene expression can be altered in cells that have properties desirable for in vitro protein production (e.g., pharmaceuticals) or in vivo protein delivery methods (e.g. gene therapy). Figures 5 and 6 illustrate two strategies for transcriptionally activating the hEPO gene.

-42-

Using the methods and DNA constructs or plasmids taught herein or modifications thereof which are apparent to one of ordinary skill in the art, exogenous DNA which encodes a therapeutic product (e.g., protein, ribozyme, anti-sense RNA) can be inserted at preselected sites in the genome of vertebrate (e.g., mammalian, both human and nonhuman) primary or secondary cells.

The present invention will now be illustrated by the following examples, which are not intended to be limiting in any way.

-43-

EXAMPLES

EXAMPLE 1. PRODUCTION OF TRANSFECTED CELL STRAINS BY GENE TARGETING

Gene targeting occurs when transfecting DNA either
5 integrates into or partially replaces chromosomal DNA
sequences through a homologous recombinant event. While
such events can occur in the course of any given transfection
experiment, they are usually masked by a vast excess
of events in which plasmid DNA integrates by nonhomolo-
10 gous, or illegitimate, recombination.

a. GENERATION OF A CONSTRUCT USEFUL FOR SELECTION OF GENE TARGETING EVENTS IN HUMAN CELLS

One approach to selecting the targeted events is by
genetic selection for the loss of a gene function due to
15 the integration of transfecting DNA. The human HPRT locus
encodes the enzyme hypoxanthine-phosphoribosyl transfer-
ase. *hprt*⁻ cells can be selected for by growth in medium
containing the nucleoside analog 6-thioguanine (6-TG):
cells with the wild-type (HPRT⁺) allele are killed by
20 6-TG, while cells with mutant (*hprt*⁻) alleles can survive.
Cells harboring targeted events which disrupt HPRT gene
function are therefore selectable in 6-TG medium.

To construct a plasmid for targeting to the HPRT
locus, the 6.9 kb HindIII fragment extending from posi-
25 tions 11,960-18,869 in the HPRT sequence (Genebank name
HUMHPRTB; Edwards, A. et al., Genomics 6:593-608 (1990))
and including exons 2 and 3 of the HPRT gene, is subcloned
into the HindIII site of pUC12. The resulting clone is
cleaved at the unique XhoI site in exon 3 of the HPRT gene
30 fragment and the 1.1 kb SalI-XhoI fragment containing the
neo gene from pMC1Neo (Stratagene) is inserted, disrupting
the coding sequence of exon 3. One orientation, with the
direction of neo transcription opposite that of HPRT
transcription was chosen and designated pE3Neo. The

-44-

replacement of the normal HPRT exon 3 with the neo-disrupted version will result in an $hprt^-$, 6-TG resistant phenotype. Such cells will also be G418 resistant.

b. GENE TARGETING IN AN ESTABLISHED HUMAN FIBROBLAST
5 CELL LINE

As a demonstration of targeting in immortalized cell lines, and to establish that pE3Neo functions properly in gene targeting, the human fibrosarcoma cell line HT1080 (ATCC CCL 121) was transfected with pE3Neo by electroporation.
10 tion.

HT1080 cells were maintained in HAT (hypoxanthine/aminopterin/xanthine) supplemented DMEM with 15% calf serum (Hyclone) prior to electroporation. Two days before electroporation, the cells are switched to the same medium
15 without aminopterin. Exponentially growing cells were trypsinized and diluted in DMEM/15% calf serum, centrifuged, and resuspended in PBS (phosphate buffered saline) at a final cell volume of 13.3 million cells per ml. pE3Neo is digested with HindIII, separating the 8 kb
20 HPRT-neo fragment from the pUC12 backbone, purified by phenol extraction and ethanol precipitation, and resuspended at a concentration of 600 $\mu\text{g/ml}$. 50 μl (30 μg) was added to the electroporation cuvette (0.4 cm electrode gap; Bio-Rad Laboratories), along with 750 μl of the cell
25 suspension (10 million cells). Electroporation was at 450 volts, 250 μFarads (Bio-Rad Gene Pulser; Bio-Rad Laboratories). The contents of the cuvette were immediately added to DMEM with 15% calf serum to yield a cell suspension of 1 million cells per 25 ml media. 25 ml of the treated
30 cell suspension was plated onto 150 mm diameter tissue culture dishes and incubated at 37°C, 5% CO_2 . 24 hrs later, a G418 solution was added directly to the plates to yield a final concentration of 800 $\mu\text{g/ml}$ G418. Five days later the media was replaced with DMEM/15% calf serum/

-45-

800 $\mu\text{g/ml}$ G418. Nine days after electroporation, the media was replaced with DMEM/15% calf serum/800 $\mu\text{g/ml}$ G418 and 10 μM 6-thioguanine. Colonies resistant to G418 and 6-TG were picked using cloning cylinders 14-16 days after the dual selection was initiated.

The results of five representative targeting experiments in HT1080 cells are shown in Table 1.

TABLE 1

Transfection	Number of Treated Cells	Number of G418 ^r 6-TG ^r Clones
1	1×10^7	32
2	1×10^7	28
3	1×10^7	24
4	1×10^7	32
5	1×10^7	66

For transfection 5, control plates designed to determine the overall yield of G418^r colonies indicated that 33,700 G418^r colonies could be generated from the initial 1×10^7 treated cells. Thus, the ratio of targeted to non-targeted events is 66/33,700, or 1 to 510. In the five experiments combined, targeted events arise at a frequency of 3.6×10^6 , or 0.00036% of treated cells.

Restriction enzyme and Southern hybridization experiments using probes derived from the neo and HPRT genes localized the neo gene to the HPRT locus at the predicted site within HPRT exon 3.

-46-

c. GENE TARGETING IN PRIMARY AND SECONDARY HUMAN SKIN FIBROBLASTS

pE3Neo is digested with HindIII, separating the 8 kb HPRT-neo fragment from the pUC12 backbone, and purified by phenol extraction and ethanol precipitation. DNA was resuspended at 2 mg/ml. Three million secondary human foreskin fibroblasts cells in a volume of 0.5 ml were electroporated at 250 volts and 960 μ Farads, with 100 μ g of HindIII pE3Neo (50 μ l). Three separate transfections were performed, for a total of 9 million treated cells. Cells are processed and selected for G418 resistance. 500,000 cells per 150 mm culture dish were plated for G418 selection. After 10 days under selection, the culture medium is replaced with human fibroblast nutrient medium containing 400 μ g/ml G418 and 10 μ M 6-TG. Selection with the two drug combination is continued for 10 additional days. Plates are scanned microscopically to localize human fibroblast colonies resistant to both drugs. The fraction of G418^r t-TG^r colonies is 4 per 9 million treated cells. These colonies constitute 0.0001% (or 1 in a million) of all cells capable of forming colonies. Control plates designed to determine the overall yield of G418^r colonies indicated that 2,850 G418^r colonies could be generated from the initial 9×10^6 treated cells. Thus, the ratio of targeted to non-targeted events is 4/2,850, or 1 to 712. Restriction enzyme and Southern hybridization experiments using probes derived from the neo and HPRT genes were used to localize the neo gene to the HPRT locus at the predicted site within HPRT exon 3 and demonstrate that targeting had occurred in these four clonal cell strains. Colonies resistant to both drugs have also been isolated by transfecting primary cells ($1/3.0 \times 10^7$).

The results of several pE3Neo targeting experiments are summarized in Table 2. HindIII digested pE3Neo was

-47-

- either transfected directly or treated with exonuclease III to generate 5' single-stranded overhangs prior to transfection (see Example 1c). DNA preparations with single-stranded regions ranging from 175 to 930 base pairs in length were tested. Using pE3neo digested with HindIII alone, 1/799 G418-resistant colonies were identified by restriction enzyme and Southern hybridization analysis as having a targeted insertion of the *neo* gene at the HPRT locus (a total of 24 targeted clones were isolated).
- 10 Targeting was maximally stimulated (approximately 10-fold stimulation) when overhangs of 175 bp were used, with 1/80 G418^r colonies displaying restriction fragments that are diagnostic for targeting at HPRT (a total of 9 targeted clones were isolated). Thus, using the conditions and
- 15 recombinant DNA constructs described here, targeting is readily observed in normal human fibroblasts and the overall targeting frequency (the number of targeted clones divided by the total number of clones stably transfected to G418-resistance) can be stimulated by transfection with
- 20 targeting constructs containing single-stranded overhang-ing tails, by the method as described in Example 1e.

TABLE 2
TARGETING TO THE HPRT LOCUS IN HUMAN FIBROBLASTS

<u>pE3neo</u> <u>Treatment</u>	<u>Number of</u> <u>Experiments</u>	<u>Number Targeted</u> <u>Per G418^r Colony</u>	<u>Total Number of</u> <u>Targeted Clone</u>
HindIII digest	6	1/799	24
175 bp overhang	1	1/80	9
350 bp overhang	3	1/117	20
930 bp overhang	1	1/144	1

-48-

d. GENERATION OF A CONSTRUCT FOR TARGETED INSERTION OF A
GENE OF THERAPEUTIC INTEREST INTO THE HUMAN GENOME
AND ITS USE IN GENE TARGETING

5 A variant of pE3Neo, in which a gene of therapeutic
interest is inserted within the HPRT coding region, adja-
cent to or near the neo gene, can be used to target a gene
of therapeutic interest to a specific position in a recip-
ient primary or secondary cell genome. Such a variant of
pE3Neo can be constructed for targeting the hGH gene to
10 the HPRT locus.

pXGH5 (schematically presented in Figure 3) is di-
gested with EcoRI and the 4.1 kb fragment containing the
hGH gene and linked mouse metallothionein (mMT) promoter
is isolated. The EcoRI overhangs are filled in with the
15 Klenow fragment from *E. coli* DNA polymerase. Separately,
pE3Neo is digested with XhoI, which cuts at the junction
of the neo fragment and HPRT exon 3 (the 3' junction of
the insertion into exon 3). The XhoI overhanging ends of
the linearized plasmid are filled in with the Klenow
20 fragment from *E. coli* DNA polymerase, and the resulting
fragment is ligated to the 4.1 kb blunt-ended hGH-mMT
fragment. Bacterial colonies derived from the ligation
mixture are screened by restriction enzyme analysis for a
single copy insertion of the hGH-mMT fragment and one
25 orientation, the hGH gene transcribed in the same direc-
tion as the neo gene, is chosen and designated pE3Neo/hGH.
pE3Neo/hGH is digested with HindIII, releasing the 12.1 kb
fragment containing HPRT, neo and mMT-hGH sequences.
Digested DNA is treated and transfected into primary or
30 secondary human fibroblasts as described in Example 1c.
G418^r TG^r colonies are selected and analyzed for targeted
insertion of the mMT-hGH and neo sequences into the HPRT
gene as described in Example 1c. Individual colonies are
assayed for hGH expression using a commercially available
35 immunoassay (Nichols Institute).

-49-

Secondary human fibroblasts were transfected with pE3Neo/hGH and thioguanine-resistant colonies were analyzed for stable hGH expression and by restriction enzyme and Southern hybridization analysis. Of thirteen TG^r colonies analyzed, eight colonies were identified with an insertion of the hGH gene into the endogenous HPRT locus. All eight strains stably expressed significant quantities of hGH, with an average expression level of 22.7 μ g/10⁶ cells/24 hours. Alternatively, plasmid pE3neoEPO, Figure 4, may be used to target EPO to the human HPRT locus.

The use of homologous recombination to target a gene of therapeutic interest to a specific position in a cell's genomic DNA can be expanded upon and made more useful for producing products for therapeutic purposes (e.g., pharmaceuticals, gene therapy) by the insertion of a gene through which cells containing amplified copies of the gene can be selected for by exposure of the cells to an appropriate drug selection regimen. For example, pE3neo/hGH (Example 1d) can be modified by inserting the *dhfr*, *ada*, or CAD gene at a position immediately adjacent to the hGH or neo genes in pE3neo/hGH. Primary, secondary, or immortalized cells are transfected with such a plasmid and correctly targeted events are identified. These cells are further treated with increasing concentrations of drugs appropriate for the selection of cells containing amplified genes (for *dhfr*, the selective agent is methotrexate, for CAD the selective agent is N-(phosphonacetyl)-L-aspartate (PALA), and for *ada* the selective agent is an adenine nucleoside (e.g., alanosine). In this manner the integration of the gene of therapeutic interest will be coamplified along with the gene for which amplified copies are selected. Thus, the genetic engineering of cells to produce genes for therapeutic uses can be readily controlled by preselecting the site at which the targeting

-50-

construct integrates and at which the amplified copies reside in the amplified cells.

e. MODIFICATION OF DNA TERMINI TO ENHANCE TARGETING

Several lines of evidence suggest that 3'-overhanging
5 ends are involved in certain homologous recombination
pathways of E. coli, bacteriophage, S. cerevisiae and
Xenopus laevis. In Xenopus laevis oocytes, molecules with
3'-overhanging ends of several hundred base pairs in
length underwent recombination with similarly treated
10 molecules much more rapidly after microinjection than
molecules with very short overhangs (4 bp) generated by
restriction enzyme digestion. In yeast, the generation of
3'-overhanging ends several hundred base pairs in length
appears to be a rate limiting step in meiotic recombination.
15 No evidence for an involvement of 3'-overhanging ends
in recombination in human cells has been reported, and in
no case have modified DNA substrates of any sort been
shown to promote targeting (one form of homologous recombination) in any species. The experiment described in the
20 following example and Example 1c suggests that 5'-overhanging ends are effective for stimulating targeting in
primary, secondary and immortalized human fibroblasts.

There have been no reports on the enhancement of
targeting by modifying the ends of the transfecting DNA
25 molecules. This example serves to illustrate that modification of the ends of linear DNA molecules, by conversion of the molecules' termini from a double-stranded form to a single-stranded form, can stimulate targeting into the genome of primary and secondary human fibroblasts.

30 1100 µg of plasmid pE3Neo (Example 1a) is digested with HindIII. This DNA can be used directly after phenol extraction and ethanol precipitation, or the 8 kb HindIII fragment containing only HPRT and the neo gene can be separated away from the pUC12 vector sequences by gel

-51-

electrophoresis. ExoIII digestion of the HindIII digested DNA results in extensive exonucleolytic digestion at each end, initiating at each free 3' end, and leaving 5'-overhanging ends. The extent of exonucleolytic action and, hence, the length of the resulting 5'-overhangs, can be controlled by varying the time of ExoIII digestion. ExoIII digestion of 100 μ g of HindIII digested pE3Neo is carried out according to the supplier's recommended conditions, for times of 30 sec, 1 min, 1.5 min, 2 min, 2.5 min, 3 min, 3.5 min, 4 min, 4.5 min, and 5 min. To monitor the extent of digestion an aliquot from each time point, containing 1 μ g of ExoIII treated DNA, is treated with mung bean nuclease (Promega), under conditions recommended by the supplier, and the samples fractionated by gel electrophoresis. The difference in size between non-treated, HindIII digested pE3Neo and the same molecules treated with ExoIII and mung bean nuclease is measured. This size difference divided by two gives the average length of the 5'-overhang at each end of the molecule. Using the time points described above and digestion at 30°, the 5'-overhangs produced should range from 100 to 1,000 bases.

60 μ g of ExoIII treated DNA (total HindIII digest of pE3Neo) from each time point is purified and electroporated into primary, secondary, or immortalized human fibroblasts under the conditions described in Example 1c. The degree to which targeting is enhanced by each ExoIII treated preparation is quantified by counting the number of G418^r 6-TG^r colonies and comparing these numbers to targeting with HindIII digested pE3Neo that was not treated with ExoIII.

The effect of 3'-overhanging ends can also be quantified using an analogous system. In this case HindIII digested pE3Neo is treated with bacteriophage T7 gene 6 exonuclease (United States Biochemicals) for varying time

-52-

intervals under the supplier's recommended conditions. Determination of the extent of digestion (average length of 3'-overhang produced per end) and electroporation conditions are as described for ExoIII treated DNA. The degree to which targeting is enhanced by each T7 gene 6 exonuclease treated preparation is quantified by counting the number of G418^r 6-TG^r colonies and comparing these numbers to targeting with HindIII digested pE3Neo that was not treated with T7 gene 6 exonuclease.

Other methods for generating 5' and 3' overhanging ends are possible, for example, denaturation and annealing of two linear molecules that partially overlap with each other will generate a mixture of molecules, each molecule having 3'-overhangs at both ends or 5'-overhangs at both ends, as well as reannealed fragments indistinguishable from the starting linear molecules. The length of the overhangs is determined by the length of DNA that is not in common between the two DNA fragments.

f. CONSTRUCTION OF TARGETING PLASMIDS FOR PLACING THE HUMAN ERYTHROPOIETIN GENE UNDER THE CONTROL OF THE MOUSE METALLOTHIONEIN PROMOTER IN PRIMARY, SECONDARY AND IMMORTALIZED HUMAN FIBROBLASTS

The following serves to illustrate one embodiment of the present invention, in which the normal positive and negative regulatory sequences upstream of the human erythropoietin (hEPO) gene are altered to allow expression of human erythropoietin in primary, secondary or immortalized human fibroblasts, which do not express hEPO in significant quantities as obtained.

A region lying exclusively upstream of the human EPO coding region can be amplified by PCR. Three sets of primers useful for this purpose were designed after analysis of the published human EPO sequence [Genbank designa-

-53-

tion HUMERPA; Lin, F-K., et al., Proc. Natl. Acad. Sci., USA 82:7580-7584 (1985)]. These primer pairs can amplify fragments of 609, 603, or 590 bp.

TABLE 3

Primer	HUMERPA Coordinate	Sequence	Fragment Size
F1	2 → 20	5' AGCTTCTGGGCTTCCAGAC (SEQ ID NO 1)	
R2	610 → 595	5' GGGGTCCCTCAGCGAC (SEQ ID NO 2)	609 bp
F2	8 → 24	5' TGGGCTTCCAGACCCAG (SEQ ID NO 3)	
R2	610 → 595	5' GGGGTCCCTCAGCGAC	603 bp
F3	21 → 40	5' CCAGCTACTTTGCGGAACTC (SEQ ID NO 4)	
R2	610 → 595	5' GGGGTCCCTCAGCGAC	590 bp

The three fragments overlap substantially and are interchangeable for the present purposes. The 609 bp fragment, extending from -623 to -14 relative to the translation start site (HUMERPA nucleotide positions 2 to 610), is ligated at both ends with *Cla*I linkers. The resulting *Cla*I-linked fragment is digested with *Cla*I and inserted into the *Cla*I site of pBluescriptIISK/+ (Stratagene), with the orientation such that HUMERPA nucleotide position 610 is adjacent to the *Sal*I site in the plasmid polylinker). This plasmid, p5'EPO, can be cleaved, separately, at the unique *Fsp*I or *Sfi*I sites in the human EPO upstream fragment (HUMERPA nucleotide positions 150 and

-54-

405, respectively) and ligated to the mouse metallothionein promoter. Typically, the 1.8 kb EcoRI-BglII from the mMT-I gene [containing no mMT coding sequences; Hamer, D.H. and Walling M., J. Mol. Appl. Gen. 1:273 288 (1982);
5 this fragment can also be isolated by known methods from mouse genomic DNA using PCR primers designed from analysis of mMT sequences available from Genbank; i.e., MUSMTI, MUSMTIP, MUSMTIPRM] is made blunt-ended by known methods and ligated with SfiI digested (also made blunt-ended) or
10 FspI digested p5'EPO. The orientations of resulting clones are analyzed and those in which the former mMT BglII site is proximal to the SalI site in the plasmid polylinker are used for targeting primary and secondary human fibro-
15 blasts. This orientation directs mMT transcription towards HUMERPA nucleotide position 610 in the final construct. The resulting plasmids are designated p5'EPO-mMTF and p5'EPO-mMTS for the mMT insertions in the FspI and SfiI sites, respectively.

Additional upstream sequences are useful in cases
20 where it is desirable to modify, delete and/or replace negative regulatory elements or enhancers that lie upstream of the initial target sequence. In the case of EPO, a negative regulatory element that inhibits EPO expression in extrahepatic and extrarenal tissues [Semenza, G.L. et al., Mol. Cell. Biol. 10:930-938 (1990)] can
25 be deleted. A series of deletions within the 6 kb fragment are prepared. The deleted regions can be replaced with an enhancer with broad host-cell activity [e.g. an enhancer from the Cytomegalovirus (CMV)].

30 The orientation of the 609 bp 5'EPO fragment in the pBluescriptIISK/+ vector was chosen since the HUMERPA sequences are preceded on their 5' end by a BamHI (distal) and HindIII site (proximal). Thus, a 6 kb BamHI-HindIII fragment normally lying upstream of the 609 bp fragment
35 [Semenza, G. L. et al., Mol. Cell. Biol. 10:930-938

-55-

(1990)] can be isolated from genomic DNA by known methods. For example, a bacteriophage, cosmid, or yeast artificial chromosome library could be screened with the 609 bp PCR amplified fragment as a probe. The desired clone will

5 have a 6 kb BamHI-HindIII fragment and its identity can be confirmed by comparing its restriction map from a restriction map around the human EPO gene determined by known methods. Alternatively, constructing a restriction map of the human genome upstream of the EPO gene using the 609 bp

10 fragment as a probe can identify enzymes which generate a fragment originating between HUMERPA coordinates 2 and 609 and extending past the upstream BamHI site; this fragment can be isolated by gel electrophoresis from the appropriate digest of human genomic DNA and ligated into a bacterial or yeast cloning vector. The correct clone will

15 hybridize to the 609 bp 5'EPO probe and contain a 6 kb BamHI-HindIII fragment. The isolated 6 kb fragment is inserted in the proper orientation into p5'EPO, p5'EPO-mMTF, or p5'EPO-mMTS (such that the HindIII site is adjacent to HUMERPA nucleotide position 2). Additional upstream sequences can be isolated by known methods, using chromosome walking techniques or by isolation of yeast artificial chromosomes hybridizing to the 609 bp 5'EPO

20 probe.

25 The cloning strategies described above allow sequences upstream of EPO to be modified in vitro for subsequent targeted transfection of primary, secondary or immortalized human fibroblasts. The strategies describe simple insertions of the mMT promoter, as well as deletion of the

30 negative regulatory region, and deletion of the negative regulatory region and replacement with an enhancer with broad host-cell activity.

-56-

g. ACTIVATING THE HUMAN EPO GENE AND ISOLATION OF TARGETED PRIMARY, SECONDARY AND IMMORTALIZED HUMAN FIBROBLASTS BY SCREENING

For targeting, the plasmids are cut with restriction enzymes that free the insert away from the plasmid backbone. In the case of p5'EPO-mMTS, HindIII and SalI digestion releases a targeting fragment of 2.4 kb, comprised of the 1.8 kb mMT promoter flanked on the 5' and 3' sides by 405 bp and 204 base pairs, respectively, of DNA for targeting this construct to the regulatory region of the human EPO gene. This DNA or the 2.4 kb targeting fragment alone is purified by phenol extraction and ethanol precipitation and transfected into primary or secondary human fibroblasts under the conditions described in Example 1c. Transfected cells are plated onto 150 mm dishes in human fibroblast nutrient medium. 48 hours later the cells are plated into 24 well dishes at a density of 10,000 cells/cm² [approximately 20,000 cells per well; if targeting occurs at a rate of 1 event per 10⁶ clonable cells (Example 1c, then about 50 wells would need to be assayed to isolate a single expressing colony]. Cells in which the transfecting DNA has targeted to the homologous region upstream of the human EPO gene will express hEPO under the control of the mMT promoter. After 10 days, whole well supernatants are assayed for EPO expression using a commercially available immunoassay kit (Amgen). Clones from wells displaying hEPO synthesis are isolated using known methods, typically by assaying fractions of the heterogeneous populations of cells separated into individual wells or plates, assaying fractions of these positive wells, and repeating as needed, ultimately isolating the targeted colony by screening 96-well microtiter plates seeded at one cell per well. DNA from entire plate lysates can also be analyzed by PCR for amplification of a fragment using a mMT specific primer in conjunction with a primer lying

-57-

upstream of HUMERPA nucleotide position 1. This primer pair should amplify a DNA fragment of a size precisely predicted based on the DNA sequence. Positive plates are trypsinized and replated at successively lower dilutions, and the DNA preparation and PCR steps repeated as needed to isolate targeted cells.

The targeting schemes herein described can also be used to activate hGH expression in immortalized human cells (for example, HT1080 cells (ATCC CCL 121), HeLa cells and derivatives of HeLa cells (ATCC CCL2, 2.1 and 2.2), MCF-7 breast cancer cells (ATCC HBT 22), K-562 leukemia cells (ATCC CCL 232), KB carcinoma cells (ATCC CCL 17), 2780AD ovarian carcinoma cells (Van der Blick, A.M. *et al.*, Cancer Res. 48:5927-5932 (1988), Raji cells (ATCC CCL 86), Jurkat cells (ATCC TIB 152), Namalwa cells (ATCC CRL 1432), HL-60 cells (ATCC CCL 240), Daudi cells (ATCC CCL 213), RPMI 8226 cells (ATCC CCL 155), U-937 cells (ATCC CRL 1593), Bowes Melanoma cells (ATCC CRL 9607), WI-38VA13 subline 2R4 cells (ATCC CLL 75.1), MOLT-4 cells (ATCC CRL 1582), and various heterohybridoma cells) for the purposes of producing hGH for conventional pharmaceutical delivery.

h. ACTIVATING THE HUMAN EPO GENE AND ISOLATION OF TARGETED PRIMARY, SECONDARY AND IMMORTALIZED HUMAN FIBROBLASTS BY A POSITIVE OR A COMBINED POSITIVE/NEGATIVE SELECTION SYSTEM

The strategy for constructing p5'EPO-mMTF, p5'EPO-mMTS, and derivatives of such with the additional upstream 6 kb BamHI-HindIII fragment can be followed with the additional step of inserting the neo gene adjacent to the mMT promoter. In addition, a negative selection marker, for example, gpt [from pMSG (Pharmacia) or another suitable source], can be inserted adjacent to the HUMERPA sequences in the pBluescriptIISK/+ polylinker. In the former case,

-58-

G418^r colonies are isolated and screened by PCR amplification or restriction enzyme and Southern hybridization analysis of DNA prepared from pools of colonies to identify targeted colonies. In the latter case, G418^r colonies are placed in medium containing 6-thioxanthine to select against the integration of the gpt gene [Besnard, C. *et al.*, Mol. Cell. Biol. 7:4139-4141 (1987)]. In addition, the HSV-TK gene can be placed on the opposite side of the insert as gpt, allowing selection for neo and against both gpt and TK by growing cells in human fibroblast nutrient medium containing 400 µg/ml G418, 100 µM 6-thioxanthine, and 25 µg/ml gancyclovir. The double negative selection should provide a nearly absolute selection for true targeted events and Southern blot analysis provides an ultimate confirmation.

The targeting schemes herein described can also be used to activate hEPO expression in immortalized human cells (for example, HT1080 cells (ATCC CCL 121), HeLa cells and derivatives of HeLa cells (ATCC CCL2, 2.1 and 2.2), MCF-7 breast cancer cells (ATCC HBT 22), K-562 leukemia cells (ATCC CCL 232), KB carcinoma cells (ATCC CCL 17), 2780AD ovarian carcinoma cells (Van der Blick, A.M. *et al.*, Cancer Res. 48:5927-5932 (1988), Raji cells (ATCC CCL 86), Jurkat cells (ATCC TIB 152), Namalwa cells (ATCC CRL 1432), HL-60 cells (ATCC CCL 240), Daudi cells (ATCC CCL 213), RPMI 8226 cells (ATCC CCL 155), U-937 cells (ATCC CRL 1593), Bowes Melanoma cells (ATCC CRL 9607), WI-38VA13 subline 2R4 cells (ATCC CLL 75.1), MOLT-4 cells (ATCC CRL 1582), and various heterohybridoma cells) for the purposes of producing hEPO for conventional pharmaceutical delivery.

-59-

i. CONSTRUCTION OF TARGETING PLASMIDS FOR PLACING THE HUMAN GROWTH HORMONE GENE UNDER THE CONTROL OF THE MOUSE METALLOTHIONEIN PROMOTER IN PRIMARY, SECONDARY OR IMMORTALIZED HUMAN FIBROBLASTS

5 The following example serves to illustrate one embodiment of the present invention, in which the normal regulatory sequences upstream of the human growth hormone gene are altered to allow expression of human growth hormone in primary, secondary or immortalized human fibro-
10 blasts.

Targeting molecules similar to those described in Example 1f for targeting to the EPO gene regulatory region are generated using cloned DNA fragments derived from the 5' end of the human growth hormone N gene. An approxi-
15 mately 1.8 kb fragment spanning HUMGHCSA (Genbank Entry) nucleotide positions 3787-5432 (the positions of two EcoNI sites which generate a convenient sized fragment for cloning or for diagnostic digestion of subclones involving this fragment) is amplified by PCR primers designed by
20 analysis of the HUMGHCSA sequence in this region. This region extends from the middle of hGH gene N intron 1 to an upstream position approximately 1.4 kb 5' to the translational start site. pUC12 is digested with EcoRI and BamHI, treated with Klenow to generate blunt ends, and
25 recircularized under dilute conditions, resulting in plasmids which have lost the EcoRI and BamHI sites. This plasmid is designated pUC12XEB. HindIII linkers are ligated onto the amplified hGH fragment and the resulting fragment is digested with HindIII and ligated to HindIII
30 digested pUC12XEB. The resulting plasmid, pUC12XEB-5'hGH, is digested with EcoRI and BamHI, to remove a 0.5 kb fragment lying immediately upstream of the hGH transcriptional initiation site. The digested DNA is ligated to the 1.8 kb EcoRI-BglII from the mMT-I gene [containing no
35 mMT coding sequences; Hamer, D.H. and Walling, M., J. Mol.

-60-

Appl. Gen. 1:273-288 (1982); the fragment can also be isolated by known methods from mouse genomic DNA using PCR primers designed from analysis of mMT sequences available from Genbank; i.e., MUSMTI, MUSMTIP, MUSMTIPRM]. This
5 plasmid p5'hGH-mMT has the mMT promoter flanked on both sides by upstream hGH sequences.

The cloning strategies described above allow sequences upstream of hGH to be modified in vitro for subsequent targeted transfection of primary, secondary or immortal-
10 ized human fibroblasts. The strategy described a simple insertion of the mMT promoter. Other strategies can be envisioned, for example, in which an enhancer with broad host-cell specificity is inserted upstream of the inserted mMT sequence.

15 j. ACTIVATING THE HUMAN hGH GENE AND ISOLATION OF TARGETED PRIMARY, SECONDARY AND IMMORTALIZED HUMAN FIBROBLASTS BY SCREENING

For targeting, the plasmids are cut with restriction enzymes that free the insert away from the plasmid back-
20 bone. In the case of p5'hGH-mMT, HindIII digestion releases a targeting fragment of 2.9 kb, comprised of the 1.8 kb mMT promoter flanked on the 5' end 3' sides by DNA for targeting this construct to the regulatory region of the hGH gene. This DNA or the 2.9 kb targeting fragment
25 alone is purified by phenol extraction and ethanol precipitation and transfected into primary or secondary human fibroblasts under the conditions described in Example 11. Transfected cells are plated onto 150 mm dishes in human fibroblast nutrient medium. 48 hours later the cells are
30 plated into 24 well dishes at a density of 10,000 cells/cm² [approximately 20,000 cells per well; if targeting occurs at a rate of 1 event per 10⁶ clonable cells (Example 1c), then about 50 wells would need to be assayed to isolate a single expressing colony]. Cells in which the

-61-

transfecting DNA has targeted to the homologous region upstream of hGH will express hGH under the control of the mMT promoter. After 10 days, whole well supernatants are assayed for hGH expression using a commercially available immunoassay kit (Nichols). Clones from wells displaying hGH synthesis are isolated using known methods, typically by assaying fractions of the heterogenous populations of cells separated into individual wells or plates, assaying fractions of these positive wells, and repeating as needed, ultimately isolated the targeted colony by screening 96-well microtiter plates seeded at one cell per well. DNA from entire plate lysates can also be analyzed by PCR for amplification of a fragment using a mMT specific primer in conjunction with a primer lying downstream of HUMGHCSA nucleotide position 5,432. This primer pair should amplify a DNA fragment of a size precisely predicted based on the DNA sequence. Positive plates are trypsinized and replated at successively lower dilutions, and the DNA preparation and PCR steps repeated as needed to isolate targeted cells.

The targeting schemes herein described can also be used to activate hGH expression in immortalized human cells (for example, HT1080 cells (ATCC CCL 121), HeLa cells and derivatives of HeLa cells (ATCC CCL2, 2.1 and 2.2), MCF-7 breast cancer cells (ATCC HBT 22), K-562 leukemia cells (ATCC CCL 232), KB carcinoma cells (ATCC CCL 17), 2780AD ovarian carcinoma cells (Van der Blick, A.M. et al., Cancer Res. 48:5927-5932 (1988), Raji cells (ATCC CCL 86), Jurkat cells (ATCC TIB 152), Namalwa cells (ATCC CRL 1432), HL-60 cells (ATCC CCL 240), Daudi cells (ATCC CCL 213), RPMI 8226 cells (ATCC CCL 155), U-937 cells (ATCC CRL 1593), Bowes Melanoma cells (ATCC CRL 9607), WI-38VA13 subline 2R4 cells (ATCC CCL 75.1), MOLT-4 cells (ATCC CRL 1582), and various heterohybridoma cells)

-62-

for the purposes of producing hGH for conventional pharmaceutical delivery.

k. ACTIVATING THE HUMAN hGH GENE AND ISOLATION OF TARGETED PRIMARY, SECONDARY AND IMMORTALIZED HUMAN FIBROBLASTS BY A POSITIVE OR A COMBINED POSITIVE/
5 NEGATIVE SELECTION SYSTEM

The strategy for constructing p5'hGH-mMT can be followed with the additional step of inserting the neo gene adjacent to the mMT promoter. In addition, a negative selection marker, for example, gpt [from pMSG (Pharmacia) or another suitable source], can be inserted adjacent to the HUMGHCSA sequences in the pUC12 poly-linker. In the former case, G418^r colonies are isolated and screened by PCR amplification or restriction enzyme and Southern hybridization analysis of DNA prepared from pools of colonies to identify targeted colonies. In the latter case, G418^r colonies are placed in medium containing thioxanthine to select against the integration of the gpt gene (Besnard, C. et al., Mol. Cell. Biol. 7: 4139-4141 (1987)). In addition, the HSV-TK gene can be placed on the opposite side of the insert as gpt, allowing selection for neo and against both gpt and TK by growing cells in human fibroblast nutrient medium containing 400 µg/ml G418, 100 µM 6-thioxanthine, and 25 µg/ml gancyclovir. The double negative selection should provide a nearly absolute selection for true targeted events. Southern hybridization analysis is confirmatory.

The targeting schemes herein described can also be used to activate hGH expression in immortalized human cells (for example, HT1080 cells (ATCC CCL 121), HeLa cells and derivatives of HeLa cells (ATCC CCL2, 2.1 and 2.2), MCF-7 breast cancer cells (ATCC HBT 22), K-562 leukemia cells (ATCC CCL 232), KB carcinoma cells (ATCC CCL 17), 2780AD ovarian carcinoma cells (Van der Blick,

-63-

A.M. et al., Cancer Res. 48:5927-5932 (1988), Raji cells (ATCC CCL 86), Jurkat cells (ATCC TIB 152), Namalwa cells (ATCC CRL 1432), HL-60 cells (ATCC CCL 240), Daudi cells (ATCC CCL 213), RPMI 8226 cells (ATCC CCL 155), U-937 cells (ATCC CRL 1593), Bowes Melanoma cells (ATCC CRL 9607), WI-38VA13 subline 2R4 cells (ATCC CLL 75.1), MOLT-4 cells (ATCC CRL 1582), and various heterohybridoma cells) for the purposes of producing hGH for conventional pharmaceutical delivery.

The targeting constructs described in Examples 1f and 1i, and used in Examples 1g, 1h, 1j and 1k can be modified to include an amplifiable selectable marker (e.g., ada, dhfr, or CAD) which is useful for selecting cells in which the activated endogenous gene, and the amplifiable selectable marker, are amplified. Such cells, expressing or capable of expressing the endogenous gene encoding a therapeutic product can be used to produce proteins (e.g., hGH and hEPO) for conventional pharmaceutical delivery or for gene therapy.

1. TRANSFECTION OF PRIMARY AND SECONDARY FIBROBLASTS WITH EXOGENOUS DNA AND A SELECTABLE MARKER GENE BY ELECTROPORATION

Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na_2HPO_4 , 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension

-64-

contains approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

Supercoiled plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad.) The final DNA
5 concentration is generally at least 120 $\mu\text{g/ml}$. 0.5 ml of the cell suspension (containing approximately 1.5×10^6 cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad).
10 Capacitance and voltage are set at 960 μF and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of
15 approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of pre-
20 warmed nutrient media (as above with 15% calf serum) in a 10 cm dish and incubated as described above. The following day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours. Subculture of cells to determine cloning efficiency and to
25 select for G418-resistant colonies is performed the following day. Cells are trypsinized, counted and plated; typically, fibroblasts are plated at 10^3 cells/10 cm dish for the determination of cloning efficiency and at $1-2 \times 10^4$ cells/10 cm dish for G418 selection.

30 Human fibroblasts are selected for G418 resistance in medium consisting of 300-400 $\mu\text{g/ml}$ G418 (Geneticin, disulfate salt with a potency of approximately 50%; Gibco) in fibroblasts nutrient media (with 15% calf serum). Cloning efficiency is determined in the absence of G418. The
35 plated cells are incubated for 12-14 days, at which time

-65-

colonies are fixed with formalin, stained with crystal violet and counted (for cloning efficiency plated) or isolated using cloning cylinders (for G418 plates).

Electroporation and selection of rabbit fibroblasts is performed essentially as described for human fibroblasts, with the exception of the selection conditions used. Rabbit fibroblasts are selected for G418 resistance in medium containing 1 gm/ml G418.

Fibroblasts were isolated from freshly excised human foreskins. Cultures were seeded at 50,000 cells/cm in DMEM + 10% calf serum. When cultures became confluent, fibroblasts were harvested by trypsinization and transfected by electroporation. Electroporation conditions were evaluated by transfection with the plasmid pCDNEO (Figure 5). A representative electroporation experiment using near optimal conditions (60 μ g of plasmid pCDNEO at an electroporation voltage of 250 volts and a capacitance setting of 960 μ Farads) resulted in one G418 colony per 588 treated cells (0.17% of all cells treated), or one G418 colony per 71 clonable cells (1.4%).

When nine separate electroporation experiments at near optimal conditions (60 μ g of plasmid pCDNEO at an electroporation voltage of 300 volts and a capacitance setting of 960 μ Farads) were performed, an average of one G418 colony per 1,899 treated cells (0.05%) was observed, with a range of 1/882 to 1/7,500 treated cells. This corresponds to an average of one G418 colony per 38 clonable cells (2.6%).

Low passage primary human fibroblasts were converted to hGH expressing cells by co-transfection with plasmids; pCDNEO and pXGH5. Typically, 60 μ g of an equimolar mixture of the two plasmids were transfected at near optimal conditions (electroporation voltage of 300 volts and a capacitance setting of 960 μ Farads). The results of such

-66-

an experiment resulted in one G418 colony per 14,705 treated cells.

hGH expression data for these and other cells isolated under identical transfection conditions are summarized below. Ultimately, 98% of all G418^r colonies could be expanded to generate mass cultures.

Number of G418 ^r Clones		
	Analyzed	154
Number of G418 ^r /hGH		
10	Expressing Clones	65
Average hGH Expression		
	Level	2.3 µg hGH/10 ⁶ Cells/24 hr
Maximum hGH Expression		
	Level	23.0 µg hGH/10 ⁶ Cells/24 hr

Stable transfectants also have been generated by electroporation of primary or secondary human fibroblasts with pXGH301, a DNA construct in which the neo and hGH genes are present on the same plasmid molecule. pXGH301 was constructed by a two-step procedure. The Sall-ClaI fragment from pBR322 (positions 23-651 in pBR322) was isolated and inserted into Sall-ClaI digested pCDNEO, introducing a BamHI site upstream of the SV40 early promoter region of pCDNEO. This plasmid, pBNEO was digested with BamHI and the 2.1 kb fragment containing the neo gene under the control of the SV40 early promoter, was isolated and inserted into BamHI digested pXGH5. A plasmid with a single insertion of the 2.1 kb BamHI fragment was isolated in which neo and hGH are transcribed in the same direction relative to each other. This plasmid was designated pXGH301. For example, 1.5 x 10⁶ cells were electroporated with 60 µg pXGH301 at 300 volts and 960 µFarads. G418 resistant colonies were isolated from transfected secondary fibroblasts at a frequency of 652 G418 resistant

-67-

colonies per 1.5×10^6 treated cells (1 per 2299 treated cells). Approximately 59% of these colonies express hGH.

EXAMPLE 2. CONSTRUCTION OF TARGETING PLASMIDS WHICH RESULT IN CHIMERIC TRANSCRIPTION UNITS IN WHICH HUMAN GROWTH HORMONE AND ERYTHROPOIETIN SEQUENCES ARE FUSED

The following serves to illustrate two further embodiments of the present invention, in which the normal regulatory sequences upstream of the human EPO gene are altered to allow expression of hEPO in primary or secondary fibroblast strains which do not express hEPO in detectable quantities in their untransfected state as obtained. In these embodiments, the products of the targeting events are chimeric transcription units in which the first exon of the human growth hormone gene is positioned upstream of hEPO exons 2-5. The product of transcription, splicing and translation is a protein in which amino acids 1-4 of the hEPO signal peptide are replaced with amino acid residues 1-3 of hGH. The two embodiments differ with respect to both the relative positions of the foreign regulatory sequences that are inserted and the specific pattern of splicing that needs to occur to produce the final, processed transcript.

Plasmid pXEPO-10 is designed to replace exon 1 of hEPO with exon 1 of hGH by gene targeting to the endogenous hEPO gene on human chromosome 7. Plasmid pXEPO-10 is constructed as follows. First, the intermediate plasmid pT163 is constructed by inserting the 6 kb HindIII-BamHI fragment (see Example 1f) lying upstream of the hEPO coding region into HindIII-BamHI digested pBluescriptII SK+ (Stratagene, LaJolla, CA). The product of this ligation is digested with XhoI and HindIII and ligated to the 1.1 kb HindIII-XhoI fragment from pMClneoPolyA [Thomas, K. R. and Capecchi, M. R. Cell 51: 503-512 (1987) available

- 68 -

from Strategene, LaJolla, CA] to create pT163. Oligo-nucleotides 13.1 - 13.4 are utilized in polymerase chain reactions to generate a fusion fragment in which the mouse metallothionein 1 (mMT-I) promoter - hGH exon 1 sequences are additionally fused to hEPO intron 1 sequences. First, oligonucleotides 13.1 and 13.3 are used to amplify the approximately 0.73 kb mMT-I promoter - hGH exon 1 fragment from pXGH5 (Figure 5). Next, oligonucleotides 13.2 and 13.4 are used to amplify the approximately 0.57 kb fragment comprised predominantly of hEPO intron 1 from human genomic DNA. Finally, the two amplified fragments are mixed and further amplified with oligonucleotides 13.1 and 13.4 to generate the final fusion fragment (fusion fragment 3) flanked by a SalI site at the 5' side of the mMT-I moiety and an XhoI site at the 3' side of the hEPO intron 1 sequence. Fusion fragment 3 is digested with XhoI and SalI and ligated to XhoI digested pT163. The ligation mixture is transformed into *E. coli* and a clone containing a single insert of fusion fragment 3 in which the XhoI site is regenerated at the 3' side of hEPO intron 1 sequences is identified and designated pXEPO-10.

```

13.1      5'  TTTTGTCGAC  GGTACCTTGG  TTTTTAAAC  C
              Sall   KpnI
              (SEQ ID NO 5)

25  13.2      5'  CCTAGCGGCA  ATGGCTACAG  GTGAGTACTC  GCGGGCTGGG  CG
              (SEQ ID NO 6)

      13.3      5'  CGCCAGCCC  GCGAGTACTC  ACCTGTAGCC  ATTGCCGCTA  GG
              (SEQ ID NO 7)

      13.4      5'  TTTTCTCGAG  CTAGAACAGA  TAGCCAGGCT  G
              XhoI
              (SEQ ID NO 8)
30

```

The non-boldface region of oligo 13.1 is identical to the mMT-I promoter, with the natural KpnI

-69-

site as its 5' boundary. The boldface type denotes a Sall site tail to convert the 5' boundary to a Sall site. The boldface region of oligos 13.2 and 13.3 denote hGH sequences, while the non-boldface regions are intron 1 sequences from the hEPO gene. The non-boldface region of oligo 13.4 is identical to the last 25 bases of hEPO intron 1. The boldface region includes an XhoI site tail to convert the 3' boundary of the amplified fragment to an XhoI site.

Plasmid pXEPO-11 is designed to place, by gene targeting, the mMT-I promoter and exon 1 of hGH upstream of the hEPO structural gene and promoter region at the endogenous hEPO locus on human chromosome 7. Plasmid pXEPO-11 is constructed as follows. Oligonucleotides 13.1 and 13.5 - 13.7 are utilized in polymerase chain reactions to generate a fusion fragment in which the mouse metallothionein I (mMT-I) promoter - hGH exon 1 sequences are additionally fused to hEPO sequences from -1 to -630 relative to the hEPO coding region. First, oligonucleotides 13.1 and 13.6 are used to amplify the approximately 0.75 kb mMT-I promoter - hGH exon 1 fragment from pXGH5 (Figure 5). Next, oligonucleotides 13.5 and 13.7 are used to amplify, from human genomic DNA, the approximately 0.65 kb fragment comprised predominantly of hEPO sequences from -1 to -620 relative to the hEPO coding region. Both oligos 13.5 and 13.6 contain a 10 bp linker sequence located at the hGH intron 1 - hEPO promoter region, which corresponds to the natural hEPO intron 1 splice-donor site. Finally, the two amplified fragments are mixed and further amplified with oligonucleotides 13.1 and 13.7 to generate the final fusion fragment (fusion fragment 6) flanked by a Sall site at the 5' side of the mMT-I moiety and an XhoI site at the 3' side of the hEPO promoter

-70-

region. Fusion fragment 6 is digested with XhoI and Sall and ligated to XhoI digested pT163. The ligation mixture is transformed into *E. coli* and a clone containing a single insert of fusion fragment 6 in which the XhoI site is regenerated at the 3' side of hEPO promoter sequences is identified and designated pXEPO-11.

13.5 5' GACAGCTCAC CTAGCGGCAA TGGCTACAGG TGAGTACTC
AAGCTTCTGG GCTTCCAGAC CCAG (SEQ ID NO 9)
HindIII

13.6 5' CTGGGTCTGG AAGCCCAGAA GCTTGAGTAC TCACCTGTAG
HindIII
CCATTGCCGC TAGGTGAGCT GTC (SEQ ID NO 10)

13.7 5' TTTTCTCGAG CTCCGCGCCT GGCCGGGGTC CCTC
XhoI
(SEQ ID NO 11)

The boldface regions of oligos 13.5 and 13.6 denote hGH sequences. The italicized regions correspond to the first 10 base pairs of hEPO intron 1. The remainder of the oligos correspond to hEPO sequences from -620 to -597 relative to the hEPO coding region. The non-boldface region of oligo 13.7 is identical to bases -1 to -24 relative to the hEPO coding region. The boldface region includes an XhoI site tail to convert the 3' boundary of the amplified fragment to an XhoI site.

Plasmid pXEPO-10 can be used for gene targeting by digestion with BamHI and XhoI to release the 7.3 kb fragment containing the mMT-I/hGH fusion flanked on both sides by hEPO sequences. This fragment (targeting fragment 1)

-71-

contains no hEPO coding sequences, having only sequences lying between -620 and approximately -6620 upstream of the hEPO coding region and hEPO intron 1 sequences to direct targeting to the human EPO locus. Targeting fragment 1 is

5 transfected into primary or secondary human skin fibroblasts using conditions similar to those described in Example 1c. G418-resistant colonies are picked into individual wells of 96-well plates and screened for EPO expression by an ELISA assay (R&D Systems, Minneapolis

10 MN). Cells in which the transfecting DNA integrates randomly into the human genome cannot produce EPO. Cells in which the transfecting DNA has undergone homologous recombination with the endogenous hEPO intron 1 and hEPO upstream sequences contain a chimeric gene in which the

15 mMT-I promoter and non-transcribed sequences and the hGH 5' untranslated sequences and hGH exon 1 replace the normal hEPO promoter and hEPO exon 1 (see Figure 1). Non-hEPO sequences in targeting fragment 1 are joined to hEPO sequences down-stream of hEPO intron 1. The replacement

20 of the normal hEPO regulatory region with the mMT-I promoter will activate the EPO gene in fibroblasts, which do not normally express hEPO. The replacement of hEPO exon 1 with hGH exon 1 results in a protein in which the first 4 amino acids of the hEPO signal peptide are replaced with

25 amino acids 1-3 of hGH, creating a functional, chimeric signal peptide which is removed by post-translation processing from the mature protein and is secreted from the expressing cells.

Plasmid pXEPO-11 can be used for gene targeting by

30 digestion with BamHI and XhoI to release the 7.4 kb fragment containing the mMT-I/hGH fusion flanked on both sides by hEPO sequences. This fragment (targeting fragment 2) contains no hEPO coding sequences, having only sequences lying between -1 and approximately -6620 upstream of the

35 hEPO coding region to direct targeting to the human EPO

-72-

locus. Targeting fragment 2 is transfected into primary or secondary human skin fibroblasts using conditions similar to those described in Example 1g. G418-resistant colonies are picked into individual wells of 96-well plates and screened for EPO expression by an ELISA assay (R&D Systems, Minneapolis, MN). Cells in which the transfecting DNA integrates randomly into the human genome cannot produce EPO. Cells in which the transfecting DNA has undergone homologous recombination with the endogenous hEPO promoter and upstream sequences contain a chimeric gene in which the mMT-I promoter and non-transcribed sequences, hGH 5' untranslated sequences and hGH exon 1, and a 10 base pair linker comprised of the first 10 bases of hEPO intron 1 are inserted at the HindIII site lying at position -620 relative to the hEPO coding region (see Figure 2). The localization of the mMT-I promoter upstream of the normally silent hEPO promoter will direct the synthesis, in primary or secondary skin fibroblasts, of a message reading (5' to 3') non-translated metallothionein and hGH sequences, hGH exon 1, 10 bases of DNA identical to the first 10 base pairs of hEPO intron 1, and the normal hEPO promoter and hEPO exon 1 (-620 to +13 relative to the hEPO coding sequence). The 10 base pair linker sequence from hEPO intron 1 acts as a splice-donor site to fuse hGH exon 1 to the next downstream splice acceptor site, that lying immediately upstream of hEPO exon 2. Processing of the resulting transcript will therefore splice out the hEPO promoter, exon 1, and intron 1 sequences. The replacement of hEPO exon 1 with hGH exon 1 results in a protein in which the first 4 amino acids of the hEPO signal peptide are replaced with amino acids 1-3 of hGH, creating a functional, chimeric signal peptide which is removed by post-translation processing from the mature protein and is secreted from the expressing cells.

-73-

A series of constructs related to pXEPO-10 and pXEPO-11 can be constructed, using known methods. In these constructs, the relative positions of the mMT-I promoter and hGH sequences, as well as the position at which the mMT-I/hGH sequences are inserted into hEPO upstream sequences, are varied to create alternative chimeric transcription units that facilitate gene targeting, result in more efficient expression of the fusion transcripts, or have other desirable properties. Such constructs will give similar results, such that an hGH-hEPO fusion gene is placed under the control of an exogenous promoter by gene targeting to the normal hEPO locus. For example, the 6 kb HindIII-BamHI fragment upstream of the hEPO gene (See Example 1f) has numerous restriction enzyme recognition sequences that can be utilized as sites for insertion of the neo gene and the mMT-I promoter/hGH fusion fragment. One such site, a BglII site lying approximately 1.3 kb upstream of the HindIII site, is unique in this region and can be used for insertion of one or more selectable markers and a regulatory region derived from another gene that will serve to activate hEPO expression in primary, secondary, or immortalized human cells.

First, the intermediate plasmid pT164 is constructed by inserting the 6 kb HindIII-BamHI fragment (Example 1f) lying upstream of the hEPO coding region into HindIII-BamHI digested pBluescriptII SK+ (Stratagene, LaJolla, CA). Plasmid pMC1neoPolyA [Thomas, K.R. and Capecchi, M.R. Cell 51:503-512 (1987); available from Stratagene, LaJolla, CA] is digested with BamHI and XhoI, made blunt-ended by treatment with the Klenow fragment of *E. coli* DNA polymerase, and the resulting 1.1 kb fragment is purified. pT164 is digested with BglII and made blunt-ended by treatment with the Klenow fragment of *E. coli* DNA polymerase. The two preceding blunt-ended fragments are ligated together and transformed into competent *E. coli*. Clones

-74-

with a single insert of the 1.1 kb neo fragment are isolated and analyzed by restriction enzyme analysis to identify those in which the BglII site recreated by the fusion of the blunt XhoI and BglII sites is localized 1.3 kb away from the unique HindIII site present in plasmid pT164. The resulting plasmid, pT165, can now be cleaved at the unique BglII site flanking the 5' side of the neo transcription unit.

Oligonucleotides 13.8 and 13.9 are utilized in polymerase chain reactions to generate a fragment in which the mouse metallothionein I (mMT-I) promoter - hGH exon 1 sequences are additionally fused to a 10 base pair fragment comprising a splice-donor site. The splice-donor site chosen corresponds to the natural hEPO intron 1 splice-donor site, although a larger number of splice-donor sites or consensus splice-donor sites can be used. The oligonucleotides (13.8 and 13.9) are used to amplify the approximately 0.73 kb mMT-I promoter - hGH exon 1 fragment from pXGH5 (Figure 5). The amplified fragment (fragment 7) is digested with BglII and ligated to BglII digested pT165. The ligation mixture is transformed into *E. coli* and a clone, containing a single insert of fragment 7 in which the KpnI site in the mMT-I promoter is adjacent to the 5' end of the neo gene and the mMT-I promoter is oriented such that transcription is directed towards the unique HindIII site, is identified and designated pXEPO-12.

13.8 5' **AAAAAGATCT** GGTACCTTGG TTTTAAAAAC CAGCCTGGAG
 BglII KpnI
 (SEQ ID NO 12)

The non-boldface region of oligo 13.8 is identical to the mMT-I promoter, with the natural KpnI site as its 5' boundary. The boldface type

-75-

denotes a BglII site tail to convert the 5' boundary to a BglII site.

13.9 5' TTTTAGATCT GAGTACTCAC CTGTAGCCAT TGCCGCTAGG
BgIII
(SEQ ID NO 13)

The boldface region of oligos 13.9 denote hGH sequences. The italicized region corresponds to the first 10 base pairs of hEPO intron 1. The underlined BglII site is added for plasmid construction purposes.

Plasmid pXPEO-12 can be used for gene targeting by digestion with BamHI and HindIII to release the 7.9 kb fragment containing the neo gene and the mMT-I/hGH fusion flanked on both sides by hEPO sequences. This fragment (targeting fragment 3) contains no hEPO coding sequences, having only sequences lying between approximately -620 and approximately -6620 upstream of the hEPO coding region to direct targeting upstream of the human EPO locus. Targeting fragment 3 is transfected into primary, secondary, or immortalized human skin fibroblasts using conditions similar to those described in Examples 1b and 1c. G418-resistant colonies are picked into individual wells of 96-well plates and screened for EPO expression by an ELISA assay (R&D Systems, Minneapolis MN). Cells in which the transfecting DNA integrates randomly into the human genome cannot produce hEPO. Cells in which the transfecting DNA has undergone homologous recombination with the endogenous hEPO promoter and upstream sequences contain a chimeric gene in which the mMT-I promoter and non-transcribed sequences, hGH 5' untranslated sequences, and hGH exon 1, and a 10 base pair linker comprised of the first 10 bases of hEPO intron 1 are inserted at the BglII site lying at

-76-

position approximately -1920 relative to the hEPO coding region. The localization of the mMT-I promoter upstream of the normally silent hEPO promoter will direct the synthesis, in primary, secondary, or immortalized human fibroblasts (or other human cells), of a message reading: (5' to 3') nontranslated metallothionein and hGH sequences, hGH exon 1, 10 bases of DNA identical to the first 10 base pairs of hEPO intron 1, and hEPO upstream region and hEPO exon 1 (from approximately -1920 to +13 relative to the EPO coding sequence). The 10 base pair linker sequence from hEPO intron 1 acts as a splice-donor site to fuse hGH exon 1 to a downstream splice acceptor site, that lying immediately upstream of hEPO exon 2. Processing of the resulting transcript will therefore splice out the hEPO upstream sequences, promoter region, exon 1, and intron 1 sequences. When using pXEPO-10, -11 and -12, post-transcriptional processing of the message can be improved by using in vitro mutagenesis to eliminate splice acceptor sites lying in hEPO upstream sequences between the mMT-I promoter and hEPO exon 1, which reduce level of productive splicing events needed create the desired message. The replacement of hEPO exon 1 with hGH exon 1 results in a protein in which the first 4 amino acids of the hEPO signal peptide are replaced with amino acids 1-3 of hGH, creating a functional, chimeric signal peptide which is removed by post-translation processing from the mature protein and is secreted from the expressing cells.

EXAMPLE 3. TARGETED MODIFICATION OF SEQUENCES UPSTREAM AND AMPLIFICATION OF THE TARGETED GENE

Human cells in which the hEPO gene has been activated by the methods previously described can be induced to amplify the neo/mMT-1/EPO transcription unit if the targeting plasmid contains a marker gene that can confer resistance to a high level of a cytotoxic agent by the

-77-

phenomenon of gene amplification. Selectable marker genes such as dihydrofolate reductase (*dhfr*, selective agent is methotrexate), the multifunctional CAD gene [encoding carbamyl phosphate synthase, aspartate transcarbamylase, and dihydro-orotase; selective agent is N-(phosphonoacetyl)-L-aspartate (PALA)], glutamine synthetase; selective agent is methionine sulphoximine (MSX), and adenosine deaminase (*ada*; selective agent is an adenine nucleoside), have been documented, among other genes, to be amplifiable in immortalized human cell lines (Wright, J.A. *et al.* Proc. Natl. Acad. Sci. USA 87:1791-1795 (1990); Cockett, M.I. *et al.* Bio/Technology 8:662-667 (1990)). In these studies, gene amplification has been documented to occur in a number of immortalized human cell lines. HT1080, HeLa, MCF-7 breast cancer cells, K-562 leukemia cells, KB carcinoma cells, or 2780AD ovarian carcinoma cells, among other cells, display amplification under appropriate selection conditions.

Plasmids pXEPO-10 and pXEPO-11 can be modified by the insertion of a normal or mutant *dhfr* gene into the unique HindIII sites of these plasmids. After transfection of HT1080 cells with the appropriate DNA, selection for G418-resistance (conferred by the *neo* gene), and identification of cells in which the hEPO gene has been activated by gene targeting of the *neo*, *dhfr*, and mMT-1 sequences to the correct position upstream of the hEPO gene, these cells can be exposed to stepwise selection in methotrexate (MTX) in order to select for amplification of *dhfr* and co-amplification of the linked *neo*, mMT-1, and hEPO sequences (Kaufman, R.J. Technique 2:221-236 (1990)). A stepwise selection scheme in which cells are first exposed to low levels of MTX (0.01 to 0.08 μ M), followed by successive exposure to incremental increases in MTX concentrations up to 250 μ M MTX or higher is employed. Linear incremental steps of 0.04 to 0.08 μ M MTX and successive 2-fold in-

-78-

creases in MTX concentration will be effective in selecting for amplified transfected cell lines, although a variety of relatively shallow increments will also be effective. Amplification is monitored by increases in *dhfr* gene copy number and confirmed by measuring in vitro hEPO expression. By this strategy, substantial over-expression of hEPO can be attained by targeted modification of sequences lying completely outside of the hEPO coding region.

Constructs similar to those described (Examples 1f, 1h, 1i, 1k, 2 and 7) to activate hGH expression in human cells can also be further modified to include the *dhfr* gene for the purpose of obtaining cells that overexpress the hGH gene by gene targeting to non-coding sequences and subsequent amplification.

EXAMPLE 4. TARGETING AND ACTIVATION OF THE HUMAN EPO LOCUS IN AN IMMORTALIZED HUMAN FIBROBLAST LINE

The targeting construct pXEPO-13 was made to test the hypothesis that the endogenous hEPO gene could be activated in a human fibroblast cell. First, plasmid pT22.1 was constructed, containing 63 bp of genomic hEPO sequence upstream of the first codon of the hEPO gene fused to the mouse metallothionein-1 promoter (mMT-I). Oligonucleotides 22.1 to 22.4 were used in PCR to fuse mMT-I and hEPO sequences. The properties of these primers are as follows: 22.1 is a 21 base oligonucleotide homologous to a segment of the mMT-I promoter beginning 28 bp upstream of the mMT-I KpnI site; 22.2 and 22.3 are 58 nucleotide complementary primers which define the fusion of hEPO and mMT-I sequences such that the fusion contains 28 bp of hEPO sequence beginning 35 bases upstream of the first codon of the hEPO gene, and mMT-I sequences beginning at base 29 of oligonucleotide 22.2, comprising the natural BglII site of mMT-I and extending 30 bases into mMT-I

-79-

sequence; 22.4 is 21 nucleotides in length and is homologous to hEPO sequences beginning 725 bp downstream of the first codon of the hEPO gene. These primers were used to amplify a 1.4 kb DNA fragment comprising a fusion of mMT-I and hEPO sequences as described above. The resulting fragment was digested with KpnI (the PCR fragment contained two KpnI sites: a single natural KpnI site in the mMT-I promoter region and a single natural KpnI site in the hEPO sequence), and purified. The plasmid pXEPO1 was also digested with KpnI, releasing a 1.4 kb fragment and a 6.4 kb fragment. The 6.4 kb fragment was purified and ligated to the 1.4 kb KpnI PCR fusion fragment. The resulting construct was called pT22.1. A second intermediate, pT22.2, was constructed by ligating the approximately 6 kb HindIII-BamHI fragment lying upstream of the hEPO structural gene (see Example 1f) to BamHI and HindIII digested pBSIISK+ (Stratagene, LaJolla, CA). A third intermediate, pT22.3, was constructed by first excising a 1.1 kb XhoI/BamHI fragment from pMCINEOpolyA (Stratagene, LaJolla, CA) containing the neomycin phosphotransferase gene. The fragment was then made blunt-ended with the Klenow fragment of DNA polymerase I (New England Biolabs). This fragment was then ligated to the HincII site of pBSIISK+ (similarly made blunt with DNA polymerase I) to produce pT22.3. A fourth intermediate, pT22.4, was made by purifying a 1.1 kb XhoI/HindIII fragment comprising the neo gene from pT22.3 and ligating this fragment to XhoI and HindIII digested pT22.2. pT22.4 thus contains the neo gene adjacent to the HindIII side of the BamHI-HindIII upstream hEPO fragment. Finally, pXEPO-13 was generated by first excising a 2.0 kb EcoRI/AccI fragment from pT22.1. The EcoRI site of this fragment defines the 5' boundary of the mMT-I promoter, while the AccI site of this fragment lies within hEPO exon 5. Thus, the AccI/EcoRI fragment contains a nearly complete hEPO expression unit,

-80-

missing only a part of exon 5 and the natural polyadenylation site. This 2.0 kb EcoRI/AccI fragment was purified, made blunt-ended by treatment with the Klenow fragment of DNA polymerase I, and ligated to XhoI digested, blunt-ended, pT22.4.

HT1080 cells were transfected with PvuI-BamHI digested pXEPO-13. pXEPO-13 digested in this way generates three fragments; a 1 kb vector fragment including a portion of the *amp* gene, a 1.7 kb fragment of remaining vector sequences and an approximately 9 kb fragment containing hEPO, *neo* and MMT-I sequences. This approximately 9 kb BamHI/PvuI fragment contained the following sequences in order from the BamHI site: an approximately 5.2 kb of upstream hEPO genomic sequence, the 1.1 kb *neo* transcription unit, the 0.7 kb MMT-I promoter and the 2.0 kb fragment containing hEPO coding sequence truncated within exon 5. 45 µg of pXEPO-13 digested in this way was used in an electroporation of 12 million cells (electroporation conditions were described in Example 1b). This electroporation was repeated a total of eight times, resulting in electroporation of a total of 96 million cells. Cells were mixed with media to provide a cell density of 1 million cells per ml and 1 ml aliquots were dispensed into a total of 96, 150mm tissue culture plates (Falcon) each containing a minimum of 35 ml of DMEM/15% calf serum. The following day, the media was aspirated and replaced with fresh medium containing 0.8 mg/ml G418 (Gibco). After 10 days of incubation, the media of each plate was sampled for hEPO by ELISA analysis (R & D Systems). Six of the 96 plates contained at least 10 mU/ml hEPO. One of these plates, number 18, was selected for purification of hEPO expressing colonies. Each of the 96, 150 mm plates contained approximately 600 G418 resistant colonies (an estimated total of 57,600 G418 resistant colonies on all 96 plates). The approximately 600 colonies on plate

-81-

- number 18 were trypsinized and replated at 50 cells/ml into 364 well plates (Sterilin). After one week of incubation, single colonies were visible at approximately 10 colonies per large well of the 364 well plates (these plates are comprised of 16 small wells within each of the 24 large wells). Each well was screened for hEPO expression at this time. Two of the large wells contained media with at least 20 mU/ml hEPO. Well number A2 was found to contain 15 colonies distributed among the 16 small wells. The contents of each of these small wells were trypsinized and transferred to 16 individual wells of a 96 well plate. following 7 days of incubation the media from each of these wells was sampled for hEPO ELISA analysis. Only a single well, well number 10, contained hEPO. This cell strain was designated HT165-18A2-10 and was expanded in culture for quantitative hEPO analysis, RNA isolation and DNA isolation. Quantitative measurement of hEPO production resulted in a value of 2,500 milliunits/million cells/24 hours.
- A 0.2 kb DNA probe extending from the AccI site in hEPO exon 5 to the BglII site in the 3' untranslated region was used to probe RNA isolated from HT165-18A2-10 cells. The targeting construct, pXEPO-13, truncated at the AccI site in exon 5 does not contain these AccI/BglII sequences and, therefore, is diagnostic for targeting at the hEPO locus. Only cell strains that have recombined in a homologous manner with natural hEPO sequences would produce an hEPO mRNA containing sequence homologous to the AccI/BglII sequences. HT165-18A2-10 was found to express an mRNA of the predicted size hybridizing with the 32-P labeled AccI/BglII hEPO probe on Northern blots. Restriction enzyme and Southern blot analysis confirmed that the neo gene and mMT-I promoter were targeted to one of the two hEPO alleles in HT165-18A2-10 cells.

-82-

These results demonstrate that homologous recombination can be used to target a regulatory region to a gene that is normally silent in human fibroblasts, resulting in the functional activation of that gene.

- 5 22.1 5' CACCTAAAAT GATCTCTCTG G (SEQ ID NO 14)
- 22.2 5' CGCGCCGGGT GACCACACCG GGGGCCCTAG ATCTGGTGAA
GCTGGAGCTA CGGAGTAA (SEQ ID NO 15)
- 22.3 5' TTACTCCGTA GCTCCAGCTT CACCAGATCT AGGGCCCCCG
GTGTGGTCAC CCGGCGCG (SEQ ID NO 16)
- 10 22.4 5' GTCTCACCGT GATATTCTCG G (SEQ ID NO 17)

EXAMPLE 5. PRODUCTION OF INTRONLESS GENES

Gene targeting can also be used to produce a processed gene, devoid of introns, for transfer into yeast or bacteria for gene expression and in vitro protein production. For example, hGH can be produced in yeast by the approach described below.

Two separate targeting constructs are generated. Targeting construct 1 (TC1) includes a retroviral LTR sequence, for example the LTR from the Moloney Murine Leukemia Virus (MoMLV), a marker for selection in human cells (e.g., the neo gene from Tn5), a marker for selection in yeast (e.g., the yeast URA3 gene), a regulatory region capable of directing gene expression in yeast (e.g., the GAL4 promoter), and optionally, a sequence that, when fused to the hGH gene, will allow secretion of hGH from yeast cells (leader sequence). The vector can also include a DNA sequence that permits retroviral packaging in human cells. The construct is organized such that the above sequences are flanked, on both sides, by

-83-

hGH genomic sequences which, upon homologous recombination with genomic hGH gene N sequences, will integrate the exogenous sequences in TC1 immediately upstream of hGH gene N codon 1 (corresponding to amino acid position 1 in the mature, processed protein). The order of DNA sequences upon integration is: hGH upstream and regulatory sequences, neo gene, LTR, URA3 gene, GAL4 promoter, yeast leader sequence, hGH sequences including and downstream of amino acid 1 of the mature protein. Targeting Construct 2 (TC2) includes sequences sufficient for plasmid replication in yeast (e.g., 2-micron circle or ARS sequences), a yeast transcriptional termination sequence, a viral LTR, and a marker gene for selection in human cells (e.g., the bacterial gpt gene). The construct is organized such that the above sequences are flanked on both sides by hGH genomic sequences which, upon homologous recombination with genomic hGH gene N sequences, will integrate the exogenous sequences in TC2 immediately downstream of the hGH gene N stop codon. The order of DNA sequences upon integration is: hGH exon 5 sequences, yeast transcription termination sequences, yeast plasmid replication sequences, LTR, gpt gene, hGH 3' non-translated sequences.

Linear fragments derived from TC1 and TC2 are sequentially targeted to their respective positions flanking the hGH gene. After superinfection of these cells with helper retrovirus, LTR directed transcription through this region will result in an RNA with LTR sequences on both ends. Splicing of this RNA will generate a molecule in which the normal hGH introns are removed. Reverse transcription of the processed transcript will result in the accumulation of double-stranded DNA copies of the processed hGH fusion gene. DNA is isolated from the doubly-targeted, retrovirally-infected cells, and digested with an enzyme that cleaves the transcription unit once within the LTR. The digested material is ligated under conditions that promote

-84-

circularization, introduced into yeast cells, and the cells are subsequently exposed to selection for the URA3 gene. Only cells which have taken up the URA3 gene (linked to the sequences introduced by TC1 and TC2 and the processed hGH gene) can grow. These cells contain a plasmid which will express the hGH protein upon galactose induction and secrete the hGH protein from cells by virtue of the fused yeast leader peptide sequence which is cleaved away upon secretion to produce the mature, biologically active, hGH molecule.

Expression in bacterial cells is accomplished by simply replacing, in TC1 and TC2, the ampicillin-resistance gene from pBR322 for the yeast URA3 gene, the tac promoter (deBoer *et al.*, Proc. Natl. Acad. Sci. 80:21-25 (1983)) for the yeast GAL4 promoter, a bacterial leader sequence for the yeast leader sequence, the pBR322 origin of replication for the 2-micron circle or ARS sequence, and a bacterial transcriptional termination (e.g., trpA transcription terminator; Christie, G.E. *et al.*, Proc. Natl. Acad. Sci. 78:4180-4184 (1981)) sequence for the yeast transcriptional termination sequence. Similarly, hEPO can be expressed in yeast and bacteria by simply replacing the hGH targeting sequences with hEPO targeting sequences, such that the yeast or bacterial leader sequence is positioned immediately upstream of hEPO codon 1 (corresponding to amino acid position 1 in the mature processed protein).

EXAMPLE 6. ACTIVATION AND AMPLIFICATION OF THE EPO GENE IN AN IMMORTALIZED HUMAN CELL LINE

Incorporation of a *dhfr* expression unit into the unique HindIII site of pXEPO-13 (see Example 4) results in a new targeting vector capable of dual selection and selection of cells in which the *dhfr* gene is amplified. The single HindIII site in pXEPO-13 defines the junction

-85-

of the *neo* gene and genomic sequence naturally residing upstream of the human EPO gene. Placement of a *dhfr* gene at this site provides a construct with the *neo* and *dhfr* genes surrounded by DNA sequence derived from the natural hEPO locus. Like pXEPO-13, derivatives with the *dhfr* gene inserted are useful to target to the hEPO locus by homologous recombination. Such a construct designated pREPO4, is represented in Figure 6. The plasmid includes exons 1-4 and part of exon 5 of the human EPO gene, as well as the HindIII-BamHI fragment lying upstream of the hEPO coding region. pSVe, pTK and pmMT-I correspond to the promoters from the SV40 early region, the Herpes Simplex Virus (HSV) thymidine kinase (TK) gene and the mouse metallothionein-I gene. It was produced as follows: HindIII-digested pXEPO-13 was purified and made blunt with the Klenow fragment of DNA polymerase I. To obtain a *dhfr* expression unit, the plasmid construct pF8CIS9080 (Eaton et al., Biochemistry 25:8343-8347 (1986)) was digested with EcoRI and SalI. A 2 Kb fragment containing the *dhfr* expression unit was purified from this digest and made blunt with Klenow fragment of DNA polymerase I. This *dhfr*-containing fragment was then ligated to the blunted HindIII site of pXEPO-13. An aliquot of this ligation was transformed into E. coli and plated on ampicillin selection plates. Following an overnight incubation at 37°C, individual bacterial colonies were observed, picked and grown. Miniplasmid preparations were made from these cultures and the resulting DNA was then subjected to restriction enzyme digestion with the enzymes BglI+HindIII, and SfiI in order to determine the orientation of the inserted *dhfr* fragments. Plasmid DNA from one of these preparations was found to contain such a 2 Kb insertion of the *dhfr* fragment. The transcription orientation of the *dhfr* expression unit in this plasmid was found to be opposite that of

-86-

the adjacent *neo* gene. This is the construct designated pREPO4.

Plasmid pREPO4 was used to amplify the hEPO locus in cells subsequent to activation of the endogenous hEPO gene by homologous recombination. Gene activation with this construct allows selection for increased DHFR expression by the use of the drug methotrexate (MTX). Typically, increased DHFR expression would occur by an increase in copy number through DNA amplification. The net result would be co-amplification of the activated hEPO gene along with *dhfr* sequences. Co-amplification of the activated EPO locus should result in increased EPO expression.

Targeting experiments were performed in HT1080 cells with pREPO4. hEPO expressing line HTREPO-52 was isolated. This line was analyzed quantitatively for EPO production and by Southern and Northern blot. This strain was found to be targeted with a single copy of *dhfr/neo/mMT-1* sequences. Expression levels obtained under 0.8 mg/ml G418 selection were approximately 1300 mU/million cells/day. Because the targeted EPO locus contained a *dhfr* expression unit, it was possible to select for increased expression of DHFR with the antifolate drug, MTX. This strain was therefore subjected to stepwise selection in 0.02, 0.05, 0.1, 0.2 and 0.4 μ M MTX. Results of initial selection of this strain are shown in Table 4 and Figure 7.

-87-

TABLE 4

<u>Cell Line</u>	<u>MTX (μM)</u>	<u>mU/ Million Cells/ 24h</u>
52C20-5-0	0	1368
52C20-5-.01	0.01	1744
52C20-5-.02	0.02	11643
52C20-5-0.05	0.05	24449
52-3-5-0.10	0.1	37019
52-3-2-0.20	0.2	67867
52-3-2-0.4B	0.4	99919

Selection with elevated levels of MTX was successful in increasing hEPO expression in line HTREPO-52, with a 70-fold increase in EPO production seen in the cell line resistant to 0.4 μ M MTX. Confirmation of amplification of the hEPO locus was accomplished by Southern blot analysis in MTX-resistant cell lines, which revealed an approximately 10-fold increase in the copy number of the activated hEPO locus relative to the parental (untargeted) hEPO allele.

10 EXAMPLE 7: PRODUCTION OF AN hEPO FUSION GENE BY INSERTION OF THE CMV PROMOTER 1.8 KB UPSTREAM OF THE GENOMIC hEPO CODING REGION

Construction of targeting plasmid pREPO15:

15 pREPO15 was constructed by first fusing the CMV promoter to hGH exon 1 by PCR amplification. A 1.6 kb fragment was amplified from hGH expression construct

-88-

pXGH308, which has the CMV promoter region beginning at nucleotide 546 and ending at nucleotide 2105 of Genbank sequence HS5MIEP fused to the hGH sequences beginning at nucleotide 5225 and ending at nucleotide 7322 of Genbank sequence HUMGHCSA, using oligonucleotides 20 and 35. Oligo 20 (35 bp, SEQ ID NO: 18), hybridized to the CMV promoter at -614 relative to the cap site (in Genbank sequence HEHCMVP1), and included a SalI site at its 5' end. Oligo 35 (42 bp, SEQ ID NO: 19), annealed to the CMV promoter at +966 and the adjacent hGH exon 1, and included the first 10 base pairs of hEPO intron 1 (containing a portion of the splice-donor site) and a HindIII site at its 5' end. The resulting PCR fragment was digested with HindIII and SalI and gel-purified. Plasmid pT163 (Example 2) was digested with XhoI and HindIII and the approximately 1.1 kb fragment containing the neo expression unit was gel-purified. The 1.6 kb CMV promoter/hGH exon 1/splice-donor site fragment and the 1.2 kb neo fragment were ligated together and inserted into the HindIII site of pBSIISK+ (Stratagene, Inc.). The resulting intermediate plasmid (designated pBNCHS) contained a neo expression unit in a transcriptional orientation opposite to that of the CMV promoter/hGH exon 1/splice-donor site fragment). A second intermediate, pREPO5ΔHindIII, was constructed by first digesting pREPO5 with HindIII. This released two fragments of 1.9 kb and 8.7 kb, and the 8.7 Kb fragment containing EPO targeting sequences was gel purified and circularized by self-ligation. The resulting plasmid, pREPO5ΔHindIII, contained only non-coding genomic DNA sequences normally residing upstream of the hEPO gene. This included sequence from -5786 to -1 relative to EPO exon 1. The 2.8 kb fragment containing neo, the CMV promoter, hGH exon 1, and the splice-donor site was excised from pBNCHS with HindIII and gel-purified. This fragment was made blunt with the Klenow fragment of DNA

-89-

polymerase I (New England Biolabs, Inc.) and ligated to BglIII-digested and blunt-ended pREPO5ΔHindIII. BglIII cuts at a position -1779 bp upstream of hEPO exon 1 in pREPO5ΔHindIII. The resulting construct, pREPO15 (Figure 8), contained EPO upstream sequences from -5786 to -1779 relative to the hEPO coding region, the neo expression unit, the CMV promoter, hGH exon 1, a splice-donor site, and sequences from -1778 to -1 bp upstream of the hEPO coding region, with the various elements assembled, in the order listed, 5' to 3' relative to nucleotide sequence of the hEPO upstream region. For transfection of human cells, pREPO15 was digested with Not I and PvuI to liberate an 8.6 kb targeting fragment. The targeting fragment contained first and second targeting sequences of 4.0 kb and 1.8 kb, respectively, with homology to DNA upstream of the hEPO gene.

Cell culture, transfection, and identification of EPO expressing targeted clones:

All cells were maintained at 37°C, 5% CO₂ and 98% humidity in DMEM containing 10% calf serum (DMEM/10, HyClone Laboratories). Transfection of secondary human foreskin fibroblasts was performed by electroporating 12 x 10⁶ cells in PBS (GIBCO) with 100 μg of DNA at 250 volts and 960 μF. The treated cells were seeded at 1 x 10⁶ cells per 150 mm plate. The following day, the media was changed to DMEM/10 containing 0.8 mg/ml G418 (GIBCO). Selection proceeded for 14 days, at which time the media was sampled for EPO production. All colonies on plates exhibiting significant hEPO levels (> 5 mU/ml) as determined by an EPO ELISA (Genzyme Inc.) were isolated with sterile glass cloning cylinders (Bellco) and transferred to individual wells of a 96 well plate. Following incubation for 1-2 days, these wells were sampled for hEPO production by ELISA. Resulting hEPO-producing cell strains

-90-

were expanded in culture for freezing, nucleic acid isolation, and quantification of EPO production.

Transfection of HT1080 cells (ATCC CCL 121) was performed by treating 12×10^6 cells in PBS (GIBCO) with 45 μ g of DNA at 450 volts and 250 μ F. Growth and identification of clones occurred as for secondary human foreskin fibroblasts described above. Isolation of hEPO producing clonal cell lines occurred by limiting dilution. This was performed by first plating colonies harvested from the initial selection plates in pools of 10-15 colonies per well of a 24 well plate. hEPO producing pools were then plated at cell densities resulting in < 1 colony per well of a 96 well plate. Individual clones were expanded for further analysis as described for human foreskin fibroblasts above.

Characterization of EPO expressing clones:

pREP015 is devoid of any hEPO coding sequence. Upon targeting of the neo/CMV promoter/hGH exon 1/splice-donor fragment upstream of hEPO exon 1, hEPO expression occurs by transcriptional initiation from the CMV promoter, producing a primary transcript that includes CMV sequences, hGH exon 1 and the splice-donor site, 1.8 kb of upstream hEPO sequences, and the normal hEPO exons, introns, and 3' untranslated sequences. Splicing of this transcript would occur from the splice-donor site adjacent to hGH exon 1 to the next downstream splice-acceptor site, which is located adjacent to hEPO exon 2. Effectively, this results in a new intron consisting of genomic sequence upstream of the hEPO gene, the normal hEPO promoter, hEPO exon 1, and hEPO intron 1. In the mature transcript, hGH exon 1 would replace hEPO exon 1. hEPO exon 1 encodes only the first four and one-third amino acids of the 26 amino acid signal peptide, which is cleaved off of the precursor protein prior to secretion from the cell.

-91-

hGH exon 1 encodes the first three and one-third amino acids of the hGH signal peptide, which also is cleaved off of the precursor protein prior to secretion from the cell. Translation of the message in which hGH exon 1 replaces hEPO exon 1 would therefore result in a protein in which the signal peptide is a chimera of hGH and hEPO sequence. Removal of the signal peptide by the normal post-translational cleavage event will produce a mature hEPO molecule whose primary sequence is indistinguishable from the normal product.

Transfection of pREPO15 into human fibroblasts resulted in EPO expression by these cells. Table 5 shows the results of targeting experiments with pREPO15 in human fibroblasts and HT1080 cells. The targeting frequency in normal human fibroblasts was found to be $1/264$ G418^r colonies, and the targeting frequency with HT1080 cells was found to be $1/450$ G418^r colonies. hEPO production levels from each of these cell strains was quantified. An hEPO producer obtained from transfection of human fibroblasts was found to be secreting 7,679 mU/ 10^6 cells/ day (Table 5). An activated hEPO cell line from HT1080 cells was producing 12,582 mU/ 10^6 cells/ day (Table 5). These results indicated that activation of the hEPO locus was efficient and caused hEPO to be produced constitutively at relatively high levels. Restriction enzyme and Southern hybridization analysis was used to confirm that targeting events had occurred at the EPO locus.

Southern blot analysis of the human fibroblast and HT1080 clones that were targeted with pREPO15 was performed. Figure 9A shows the restriction map of the parental and targeted hEPO locus, and Figure 9B shows the results of restriction enzyme and Southern hybridization analysis of a targeted human fibroblast clone. BglII/EcoRI and BamHI digests revealed 5.9 and 6.6 kb fragments, respectively, as a result of a targeting event

-92-

at the hEPO locus (lanes T1). Both of these fragments resulted from the insertion of 2.7 kb of DNA containing the neo gene and CMV promoter sequences. Since only one of the two hEPO alleles were targeted, fragments of 4.3 kb (BglII/EcoRI) or 10.6 kb (BamHI) reflecting the unaltered hEPO locus were seen in these strains and in parental DNA (lanes HF). These results confirm that a homologous recombination event had occurred at the hEPO locus resulting in the production of a novel transcription unit which directed the production of human erythropoietin.

<u>Oligonucleotide</u>	<u>Sequence</u>
------------------------	-----------------

20	5' TTTTCTCGAG TCGACGACAT TGATTATTGA CTAGT
	(SEQ ID NO: 18)

35	5' TTTTAAGCTT GAGTACTCAC CTGTAGCCAT GGTGGATCCC GT
15	(SEQ ID NO: 19)

-93-

Table 5. Transfection of pREPO15 and Activation of hEPO Expression in Human Cells

Cell Type Transfected	Cells Treated	^a G418 ^r Colonies	^b Plates With EPO Expressors	hEPO Expressors per G418 ^r Colony	hEPO Expressors per Treated Cell	^c hEPO Expression (mU/10 ⁶ cells/24 hr)
Human Fibroblasts	3.3 x 10 ⁷	264	1	1/264	1/3.3 x 10 ⁷	7679
HT1080 Cells	3.1 x 10 ⁷	2700	6	1/450	1/5.2 x 10 ⁶	12,582

^a estimated by counting colonies on 2 plates, averaging the results and extrapolating to the total number of plates

^b medium from plates with G418^r colonies was sampled for EPO ELISA analysis and those exhibiting hEPO levels greater than 5 mU/ml were counted as EPO activation events

^c quantitative hEPO production was determined from human fibroblast strain, HF342-15 or HT1080 cell line, HTREPO15-1-6-6

EXAMPLE 8: PRODUCTION AND AMPLIFICATION OF AN hEPO FUSION GENE BY INSERTION OF THE CMV PROMOTER 1.8 KB UPSTREAM OF THE GENOMIC hEPO CODING REGION

Construction of targeting plasmid pREPO18:

- 5 pREPO18 (Figure 10) was constructed by insertion of a *dhfr* expression unit at the *Cla*I site located at the 5' end of the *neo* gene of pREPO15. To obtain a *dhfr* expression unit, the plasmid construct pF8CIS9080 [Eaton et al., Biochemistry 25: 8343-8347 (1986)] was digested with
- 10 *Eco*RI and *Sal*I. A 2 kb fragment containing the *dhfr* expression unit was purified from this digest and made blunt by treatment with the Klenow fragment of DNA polymerase I. A *Cla*I linker (New England Biolabs) was then ligated to the blunted *dhfr* fragment. The products of

-94-

this ligation were then digested with ClaI ligated to ClaI digested pREPO15. An aliquot of this ligation was transformed into *E. coli* and plated on ampicillin selection plates. Bacterial colonies were analyzed by restriction enzyme digestion to determine the orientation of the inserted *dhfr* fragment. One plasmid with *dhfr* in a transcriptional orientation opposite that of the *neo* gene was designated pREPO18(-). A second plasmid with *dhfr* in the same transcriptional orientation as that of the *neo* gene was designated pREPO18(+)

Cell culture, transfection, and identification of EPO expressing targeted clones:

All cells were maintained at 37°C, 5% CO₂, and 98% humidity in DMEM containing 10% calf serum (DMEM/10, HyClone Laboratories). Transfection of HT1080 cells (ATCC, CCL 121) occurred by treating 12x 10⁶ cells in PBS (GIBCO) with 45 µg of DNA at 450 volts and 250 µF. The treated cells were seeded at 1 x 10⁶ cells per 150 mm plate. The following day, the media was changed to DMEM/10 containing 0.8 mg/ml G418 (GIBCO). Selection proceeded for 14 days, at which time the media was sampled for hEPO production. Plates exhibiting significant hEPO production levels (> 5 mU/ml) as determined by an hEPO ELISA (Genzyme Inc.) were trypsinized and the cells were re-plated for clone isolation. Isolation of hEPO producing clonal cell lines occurred by limiting dilution, by first plating clones in pools of 10-15 colonies per well of a 24 well plate, and next plating cells from hEPO producing pools at cell densities resulting in less than 1 colony per well of a 96 well plate. Individual clones were expanded in culture for freezing, nucleic acid isolation and quantification of hEPO production.

-95-

Isolation of cells containing amplified *dhfr* sequences by methotrexate selection:

Targeted G418^r cell lines producing hEPO following transfection with pREPO18 were plated at various cell densities for selection in methotrexate (MTX). As new clones emerged following selection at one MTX concentration, they were assayed for hEPO production and re-plated at various cell densities in a higher concentration of MTX (usually double the previous concentration). This process was repeated until the desired hEPO production level was reached. At each step of MTX-resistance, DNA and RNA was isolated for respective southern and northern blot analysis.

Characterization of EPO expressing clones:

pREPO18, with two different orientations of *dhfr*, was transfected into HT1080 cells. Prior to transfection, pREPO18(+) and pREPO18(-) were digested with XbaI, releasing a 7.9 kb targeting fragment containing, in the following order, a 2.1 kb region of genomic DNA upstream of hEPO exon 1 (from -3891 to -1779 relative to the hEPO ATG start codon), a 2 kb region containing the *dhfr* gene, a 1.1 kb region containing the *neo* gene, a 1.5 kb region containing the CMV promoter fused to hGH exon 1, 10 bp of hEPO intron 1 (containing a splice-donor site), followed by a 1.1 kb region of genomic DNA upstream of hEPO exon 1 (from -1778 to -678 relative to the EPO ATG start codon). Transfection and targeting frequencies from two experiments are shown in Table 6. Five primary G418^r clones were isolated from these experiments. These were expanded in culture for quantitative analysis of hEPO expression (Table 7). As pREPO18 contained the *dhfr* gene, it is possible to select for cells containing amplified copies of the targeting construct using MTX as described in Example 6. G418^r clones confirmed to be targeted to the

-96-

hEPO locus by restriction enzyme and Southern hybridization analysis were subjected to stepwise selection in MTX as described.

Table 6: Targeting of pREPO18 in HT1080 cells

Construct	DNA Digest	Cells Treated	G418 ^r Colonies	Plates With hEPO Expressors	hEPO Expressors- /G418 ^r Colony	Primary Clones Analyzed
pREPO18 (-)	XbaI	36 x 10 ⁶	16,980	39	1/435	1
pREPO18 (+)	XbaI	36 x 10 ⁶	19,290	41	1/470	4

Table 7. hEPO production in HT1080 Cell lines targeted with pREPO18

Cell Line	Construct	hEPO mU/10 ⁶ Cells/24 hr
18B3-147	pREPO18 (+)	24759
18B3-181	pREPO18 (+)	20831
18B3-145	pREPO18 (+)	17586
18B3-168	pREPO18 (+)	5293
18A3-119	pREPO18 (-)	2881

-97-

EXAMPLE 9: ACTIVATION AND AMPLIFICATION OF ENDOGENOUS
 α -INTERFERON, GM-CSF, G-CSF AND FSH β GENES IN
IMMORTALIZED HUMAN CELLS

A wide variety of endogenous cellular genes can be
5 activated and amplified using the methods and DNA con-
structs of the invention. The following describes a
general strategy for activating and amplifying the human
 α -interferon (leukocyte interferon), GM-CSF (colony stimu-
lating factor-granulocyte/macrophage), G-CSF (colony
10 stimulating factor-granulocyte) and FSH β (follicle stimu-
lating hormone beta subunit) genes.

α -interferon

The human α -interferon gene (Genbank sequence
HUMIFNAA) encodes a 188 amino acid precursor protein
15 containing a 23 amino acid signal peptide. The gene
contains no introns. Figure 11 schematically illustrates
one strategy for activating the α -interferon gene. The
targeting construct is designed to include a first target-
ing sequence homologous to sequences upstream of the gene,
20 an amplifiable marker gene, a selectable marker gene, a
regulatory region, a CAP site, a splice-donor site, an
intron, a splice acceptor site, and a second targeting
sequence corresponding to sequences downstream of the
first targeting sequence. The second targeting sequence
25 should not extend further upstream than to position -107
relative to the normal start codon in order to avoid
undesired ATG start codons.

In this strategy the first and second targeting
sequences are immediately adjacent to each other in the
30 normal target gene, but this is not required (see below).
Amplifiable marker genes and selectable marker genes
suitable for selection are described herein. The amplifi-
able marker gene and selectable marker gene may be the
same gene, their positions may be reversed, and one or

-98-

both may be situated in the intron of the targeting construct. A selectable marker gene is optional and the amplifiable marker gene is only required when amplification is desired. The incorporation of a specific CAP site is optional. Optionally, exon sequences from another gene can be included 3' to the splice-acceptor site and 5' to the second targeting sequence in the targeting construct. The regulatory region, CAP site, splice-donor site, intron, and splice acceptor site can be isolated as a complete unit from the human elongation factor-1 α (EF-1 α ; Genbank sequence HUMEF1A) gene or the cytomegalovirus (CMV; Genbank sequence HEHCMVP1) immediate early region, or the components can be assembled from appropriate components isolated from different genes.

Genomic DNA corresponding to the upstream region of the α -interferon gene for use as targeting sequences and assembly of the targeting construct can be performed using recombinant DNA methods known by those skilled in the art. As described herein, a number of selectable and amplifiable markers can be used in the targeting constructs, and the activation and amplification can be effected in a large number of cell-types. Transfection of primary, secondary, or immortalized human cells and isolation of homologously recombinant cells expressing α -interferon can be accomplished using the methods described in Example 4, using an ELISA assay for human α -interferon (Biosource International, Camarillo, CA). Alternatively, homologously recombinant cells may be identified by PCR screening as described in Example 1g and 1j. The isolation of cells containing amplified copies of the amplifiable marker gene and the activated α -interferon locus is performed as described in Example 6.

In the homologously recombinant cells, an mRNA precursor is produced which includes the exogenous exon, splice-donor site, intron, splice-acceptor site, second

-99-

targeting sequence, and human α -interferon coding region and 3' untranslated sequences (Figure 11). Splicing of this message will generate a functional mRNA which can be translated to produce human α -interferon.

- 5 The size of the intron and thus the position of the regulatory region relative to the coding region of the gene may be varied to optimize the function of the regulatory region. Multiple exons may be present in the targeting construct. In addition, the second targeting sequence
10 does not need to lie immediately adjacent to or near the first targeting sequence in the normal gene, such that portions of the gene's normal upstream region are deleted upon homologous recombination.

GM-CSF

- 15 The human GM-CSF gene (Genbank sequence HUMGMCSFG) encodes a 144 amino acid precursor protein containing a 17 amino acid signal peptide. The gene contains four exons and three introns, and the N-terminal 50 amino acids of the precursor are encoded in the first exon. Figure 12
20 schematically illustrates a strategy for activating the GM-CSF gene. In this strategy the targeting construct is designed to include a first targeting sequence homologous to sequences upstream of the gene, an amplifiable marker gene, a selectable marker gene, a regulatory region, a CAP
25 site, an exon which encodes an amino acid sequence which is identical or functionally equivalent to that of the first 50 amino acids of GM-CSF, a splice-donor site, and a second targeting sequence corresponding to sequences downstream of the first targeting sequence. By this
30 strategy, homologously recombinant cells produce an mRNA precursor which corresponds to the exogenous exon and splice-donor site, the second targeting sequence, any sequences between the second targeting sequence and the start codon of the GM-CSF gene, and the exons, introns,

-100-

and 3' untranslated region of the GM-CSF gene (Figure 11). Splicing of this message results in the fusion of the exogenous exon to exon 2 of the endogenous GM-CSF gene which, when translated, will produce GM-CSF.

5 In this strategy the first and second targeting sequences are immediately adjacent in the normal target gene, but this is not required (see below). Amplifiable marker genes and selectable marker genes suitable for selection are described herein. The amplifiable marker
10 gene and selectable marker gene can be the same gene or their positions can be reversed. A selectable marker gene is optional and the amplifiable marker gene is only required when amplification is desired. The selectable marker and/or amplifiable marker can be positioned between
15 the splice-donor site and the second targeting sequence in the targeting construct. The incorporation of a specific CAP site is optional. The regulatory region, CAP site, and splice-donor site can be isolated as a complete unit from the human elongation factor-1 α (EF-1 α ; Genbank se-
20 quence HUMEF1A) gene or the cytomegalovirus (CMV; Genbank sequence HEHCMVP1) immediate early region, or the components can be assembled from an appropriate component isolated from different genes (such as the mMT-I promoter and CAP site, and exon 1 and a splice donor site from the
25 hGH or hEPO genes.

Other approaches can be employed, for example, the first and second targeting sequences can correspond to sequences in the first intron of the GM-CSF gene. Alternatively, a targeting construct similar to that described
30 for the α -interferon can be used, in which the targeting construct is designed to include a first targeting sequence homologous to sequences upstream of the GM-CSF gene, an amplifiable marker gene, a selectable marker gene, a regulatory region, a CAP site, a splice-donor
35 site, an intron, a splice acceptor site, and a second

-101-

targeting sequence corresponding to sequences downstream of the first targeting sequence.

In any case the second targeting sequence does not need to lie immediately adjacent to or near the first targeting sequence in the normal gene, such that portions of the gene's normal upstream region are deleted upon homologous recombination. In addition, multiple non-coding or coding exons can be present in the targeting construct. Genomic DNA corresponding to the upstream or intron regions of the human GM-CSF gene for use as targeting sequences and assembly of the targeting construct can be performed using recombinant DNA methods known by those skilled in the art. As described herein, a number of selectable and amplifiable markers can be used in the targeting constructs, and the activation can be effected in a large number of cell-types. Transfection of primary, secondary, or immortalized human cells and isolation of homologously recombinant cells expressing GM-CSF can be accomplished using the methods described in Example 4, using an ELISA assay for human GM-CSF (R&D Systems, Minneapolis, MN). Alternatively, homologously recombinant cells may be identified by PCR screening as described above. The isolation of cells containing amplified copies of the amplifiable marker gene and the activated GM-CSF locus is performed as described above.

G-CSF

The human G-CSF gene (Genbank sequence HUMGCSFG) encodes 204-207 amino acid precursor protein containing a 30 amino acid signal peptide. The gene contains five exons and four introns. The first exon encodes 13 amino acids of the signal peptide. Figure 13 schematically illustrates a strategy for activating the G-CSF gene. The targeting construct is designed to include a first targeting sequence homologous to sequences upstream of the gene,

-102-

an amplifiable marker gene, a selectable marker gene, a regulatory region, a CAP site, an exon which encodes an amino acid sequence which is identical or functionally equivalent to that of the first 13 amino acids of the G-CSF signal peptide, a splice-donor site, and a second targeting sequence corresponding to sequences downstream of the first targeting sequence. By this strategy, homologously recombinant cells produce an mRNA precursor which corresponds to the exogenous exon and splice-donor site, the second targeting sequence, any sequences between the second targeting sequence and the start codon of the G-CSF gene, and the exons, introns, and 3' untranslated region of the G-CSF gene (Figure 13). Splicing of this message results in the fusion of the exogenous exon to exon 2 of the endogenous G-CSF gene which, when translated, will produce G-CSF. The ability to functionally substitute the first 13 amino acids of the normal G-CSF signal peptide with those present in the exogenous exon allows one to make modifications in the signal peptide, and hence the secretory properties of the protein produced.

In this strategy the first and second targeting sequences are immediately adjacent in the normal target gene, but this is not required. The second targeting sequence does not need to lie immediately adjacent to or near the first targeting sequence in the normal gene, such that portions of the gene's normal upstream region are deleted upon homologous recombination. The amplifiable marker gene and selectable marker gene can be the same gene or their positions can be reversed. A selectable marker gene is optional and the amplifiable marker gene is only required when amplification is desired. The selectable marker and/or amplifiable marker can be positioned between the splice-donor site and the second targeting sequence in the targeting construct. The incorporation of a specific CAP site is optional. The regulatory region,

-103-

CAP site, and splice-donor site can be isolated as a complete unit from the human elongation factor-1 α (EF-1 α ; Genbank sequence HUMEF1A) gene or the cytomegalovirus (CMV; Genbank sequence HEHCMVP1) immediate early region, or the components can be assembled from an appropriate component isolated from different genes (such as the mMT-I promoter and CAP site, and exon 1 and a splice donor site from the hGH or EPO genes. Multiple exogenous exons, coding or non-coding, can be used in the targeting construct so long as an ATG start codon which, upon splicing, will be in-frame with the mature protein, is included in one of the exons.

Other approaches may be employed, for example, the first and second targeting sequences can correspond to sequences in the first intron of the G-CSF gene. Alternatively, a targeting construct similar to that described for the α -interferon can be used, in which the targeting construct is designed to include a first targeting sequence homologous to sequences upstream of the G-CSF gene, an amplifiable marker gene, a selectable marker gene, a regulatory region, a CAP site, a splice-donor site, an intron, a splice acceptor site, and a second targeting sequence corresponding to sequences downstream of the first targeting sequence.

Genomic DNA corresponding to the upstream or intron regions of the human G-CSF gene for use as targeting sequences and assembly of the targeting construct can be performed using recombinant DNA methods known by those skilled in the art. As described herein, a number of selectable and amplifiable markers can be used in the targeting constructs, and the activation can be effected in a large number of cell-types. Transfection of primary, secondary, or immortalized human cells and isolation of homologously recombinant cells expressing G-CSF can be accomplished using the methods described in Example 4,

-104-

using an ELISA assay for human G-CSF (R&D Systems, Minneapolis, MN). Alternatively, homologously recombinant cells may be identified by PCR screening as described above. The isolation of cells containing amplified copies of the amplifiable marker gene and the activated α -interferon locus is performed as described above.

FSH β

The human FSH β gene (Genbank sequence HUMFSH1) encodes a 129 amino acid precursor protein containing a 16 amino acid signal peptide. The gene contains three exons and two introns, with the first exon being a non-coding exon. The activation of FSH β can be accomplished by a number of strategies. One strategy is shown in Figure 14. In this strategy, a targeting construct is designed to include a first targeting sequence homologous to sequences upstream of the gene, an amplifiable marker gene, a selectable marker gene, a regulatory region, a CAP site, an exon, a splice-donor site, and a second targeting sequence corresponding to sequences downstream of the first targeting sequence. By this strategy, homologously recombinant cells produce an mRNA precursor which corresponds to the exogenous exon and splice-donor site, the second targeting sequence, any sequences between the second targeting sequence and the start codon of the FSH β gene, and the exons, introns, and 3' untranslated regions of the FSH β gene (Figure 14). Splicing of this message results in the fusion of the exogenous exon to exon 2 of the endogenous FSH β gene which, when translated, can produce FSH β . In this strategy the first and second targeting sequences are immediately adjacent in the normal target gene, but this is not required (see below).

Other approaches can be employed, for example, the first and second targeting sequences can correspond to sequences in the first intron of the FSH β gene. Alterna-

-105-

tively, a targeting construct similar to that described for the α -interferon can be used. In this strategy, the targeting construct is designed to include a first targeting sequence homologous to sequences upstream of the FSH β gene, an amplifiable marker gene, a selectable marker gene, a regulatory region, a CAP site, a splice-donor site, an intron, a splice acceptor site, and a second targeting sequence corresponding to sequences downstream of the first targeting sequence. The second targeting sequence should not extend further upstream than to position -40 relative to the normal FSH β transcriptional start site in order to avoid undesired ATG start codons. In the homologously recombinant cells, an mRNA precursor is produced which includes the exogenous exon, splice-donor site, intron, splice-acceptor site, second targeting sequence, and human FSH β coding exons, intron and 3' untranslated sequences. Splicing of this message will generate a functional mRNA which can be translated to produce human FSH β . The size of the intron and thus the position of the regulatory region relative to the coding region of the gene can be varied to optimize the function of the regulatory region.

In any activation strategy, the second targeting sequence does not need to lie immediately adjacent to or near the first targeting sequence in the normal gene, such that portions of the gene's normal upstream region are deleted upon homologous recombination. Furthermore, one targeting sequence can be upstream of the gene and one may be within an exon or intron of the FSH β gene.

The amplifiable marker gene and selectable marker gene can be the same gene, their positions can be reversed, and one or both can be situated in the intron of the targeting construct. Amplifiable marker genes and selectable marker genes suitable for selection are described herein. A selectable marker gene is optional and

-106-

the amplifiable marker gene is only required when amplification is desired. The incorporation of a specific CAP site is optional. Optionally, exon sequences from another gene can be included 3' to the splice-acceptor site and 5' to the second targeting sequence in the targeting construct. The regulatory region, CAP site, exon, splice-donor site, intron, and splice acceptor site can be isolated as a complete unit from the human elongation factor-1 α (EF-1 α ; Genbank sequence HUMEF1A) gene or the cytomegalovirus (CMV; Genbank sequence HEHCMVP1) immediate early region, or the components can be assembled from appropriate components isolated from different genes. In any case, the exogenous exon can be the same or different from the first exon of the normal FSH β gene, and multiple exons can be present in the targeting construct.

Genomic DNA corresponding to the upstream region of the FSH β gene for use as targeting sequences and assembly of the targeting construct can be performed using recombinant DNA methods known by those skilled in the art. As described herein, a number of selectable and amplifiable markers can be used in the targeting constructs, and the activation can be effected in a large number of cell-types. If desirable, the product of the activated FSH β gene can be produced in a cell type that expresses the human glycoprotein α -subunit, the product of which forms a heterodimer with the product of the FSH β gene. This may be a naturally occurring cell strain or line. Alternatively, the human glycoprotein α -subunit gene (Genbank sequence HUMGLYCA1) can be co-expressed with the product of the FSH β gene, with such co-expression accomplished by expression of the human glycoprotein α -subunit gene or cDNA under the control of a suitable promoter, or by activation of the human glycoprotein α -subunit gene through the methods described herein. Transfection of primary, secondary, or immortalized human cells and isola-

-107-

tion of homologously recombinant cells expressing FSH β can be accomplished using the methods described above using an ELISA assay for human FSH β (Accurate Chemical and Scientific, Westbury, NY). Alternatively, homologously recombinant cells may be identified by PCR screening as described above. The isolation of cells containing amplified copies of the amplifiable marker gene and the activated α -interferon locus is performed as described above.

Equivalents

10 Those skilled in the art will recognize, or be able to ascertain using not more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

-108-

CLAIMS

1. A DNA construct capable of altering the expression of a targeted gene when inserted into chromosomal DNA of a cell comprising:
 - 5 (a) a targeting sequence;
 - (b) a regulatory sequence;
 - (c) an exon; and
 - (d) an unpaired splice-donor site.
2. The DNA construct of Claim 1 wherein the exon comprises a CAP site.
10
3. The DNA construct of Claim 2 wherein the exon further comprises the nucleotide sequence ATG.
4. The DNA construct of Claim 3 wherein the exon further comprises encoding DNA which is in-frame with the targeted gene.
15
5. The DNA construct of Claim 4 wherein the encoding DNA of the exon is the same as the encoding DNA of the first exon of the targeted gene.
6. The DNA construct of Claim 4 wherein the encoding DNA of the exon is different from the encoding DNA of the first exon of the targeted gene.
20
7. The DNA construct of Claim 4 wherein the targeting sequence is homologous to a sequence within the targeted gene.
- 25 8. The DNA construct of Claim 4 wherein the targeting sequence is homologous to a sequence upstream of the coding region of the targeted gene.

-109-

9. The DNA construct of Claim 4 wherein the targeting sequence is homologous to a sequence upstream of the endogenous regulatory sequence of the targeted gene.
- 5 10. The DNA construct of Claim 4 wherein the construct further comprises a second targeting sequence homologous to a sequence within the targeted gene.
- 10 11. The DNA construct of Claim 4 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the coding region of the targeted gene.
12. The DNA construct of Claim 4 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the endogenous regulatory sequence of the targeted gene.
- 15 13. The DNA construct of Claim 4 wherein the targeted gene encodes a therapeutic protein.
- 20 14. The DNA construct of Claim 4 wherein the targeted gene encodes a hormone, a cytokine, an antigen, an antibody, an enzyme, a clotting factor, a transport protein, a receptor, a regulatory protein, a structural protein or a transcription factor.
- 25 15. The DNA construct of Claim 4 wherein the targeted gene encodes a protein selected from the group consisting of erythropoietin, calcitonin, growth hormone, insulin, insulinotropin, insulin-like growth factors, parathyroid hormone, β -interferon, γ -interferon, nerve growth factors, FSH β , TGF- β , tumor necrosis factor, glucagon, bone growth factor-2, bone growth factor-7, TSH- β , interleukin 1, interleukin 2,

-110-

- interleukin 3, interleukin 6, interleukin 11, interleukin 12, CSF-granulocyte, CSF-macrophage, CSF-granulocyte/ macrophage, immunoglobulins, catalytic antibodies, protein kinase C, glucocerebrosidase, superoxide dismutase, tissue plasminogen activator, urokinase, antithrombin III, DNase, α -galactosidase, tyrosine hydroxylase, blood clotting factor V, blood clotting factor VII, blood clotting factor VIII, blood clotting factor IX, blood clotting factor X, blood clotting factor XIII, apolipoprotein E or apolipoprotein A-I, globins, low density lipoprotein receptor, IL-2 receptor, IL-2 antagonists, alpha-1 antitrypsin, immune response modifiers, and soluble CD4.
- 15 16. The DNA construct of Claim 15 wherein the targeted gene encodes growth hormone, FSH β , G-CSF or GM-CSF.
17. The DNA construct of Claim 15 wherein the targeted gene encodes erythropoietin.
18. The DNA construct of Claim 17 wherein the encoding DNA of the exon is the same as the encoding DNA of the first exon of erythropoietin.
19. The DNA construct of Claim 17 wherein the encoding DNA of the exon is different from the encoding DNA of the first exon of erythropoietin.
- 20 20. The DNA construct of Claim 19 wherein the encoding DNA of the exon is the same as the encoding DNA of the first exon of human growth hormone.

-111-

21. The DNA construct of Claim 1 wherein the regulatory sequence is a promoter, an enhancer, a scaffold-attachment region or a transcription factor binding site.
- 5 22. The DNA construct of Claim 21 wherein the regulatory sequence is a promoter.
23. The DNA construct of Claim 22 further comprising an additional regulatory sequence.
- 10 24. The DNA construct of Claim 22 wherein the construct further comprises an enhancer.
25. The DNA construct of Claim 24 further comprising one or more selectable markers.
26. The DNA construct of Claim 25 further comprising an amplifiable marker gene.
- 15 27. The DNA construct of Claim 21 wherein the regulatory sequence is a regulatory sequence of the mouse metallothionein-I gene, a regulatory sequence of an SV-40 gene, a regulatory sequence of a cytomegalo-virus gene, a regulatory sequence of a collagen gene,
20 a regulatory sequence of an actin gene, a regulatory sequence of an immunoglobulin gene, a regulatory sequence of the HMG-CoA reductase gene or a regulatory sequence of the EF-1 α gene.
- 25 28. A method of making a homologously recombinant cell wherein the expression of a targeted gene is altered, comprising the steps of:
 - (a) transfecting a cell with a DNA construct, the DNA construct comprising:

-112-

- 5 (i) a targeting sequence;
(ii) a regulatory sequence;
(iii) an exon; and
(iv) an unpaired splice-donor site, thereby
producing a transfected cell; and
(b) maintaining the transfected cell under conditions appropriate for homologous recombination.
29. The method of Claim 28 wherein the exon comprises a CAP site.
- 10 30. The method of Claim 29 wherein the exon comprises the nucleotide sequence ATG.
31. The method of Claim 30 wherein the exon further comprises encoding DNA in-frame with the targeted gene.
- 15 32. The method of Claim 31 wherein the encoding DNA of the exon is the same as the encoding DNA of the first exon of erythropoietin.
33. The method of Claim 31 wherein the encoding DNA of the exon is different from the encoding DNA of the first exon of erythropoietin.
- 20 34. The method of Claim 31 wherein the targeting sequence is homologous to a sequence within the targeted gene.
35. The method of Claim 31 wherein the targeting sequence is homologous to a sequence upstream of the coding region of the targeted gene.
- 25

-113-

36. The method of Claim 31 wherein the targeting sequence is homologous to a sequence upstream of the endogenous regulatory sequence of the targeted gene.
- 5 37. The method of Claim 31 wherein the construct further comprises a second targeting sequence homologous to a sequence within the targeted gene.
- 10 38. The method of Claim 31 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the coding region of the targeted gene.
39. The method of Claim 31 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the endogenous regulatory sequence of the targeted gene.
- 15 40. The method of Claim 31 wherein the cell is a human cell.
41. The method of Claim 28 wherein the targeted gene encodes erythropoietin.
- 20 42. The method of Claim 31 wherein the encoding DNA is the same as the encoding DNA of the first exon of erythropoietin.
43. The method of Claim 31 wherein the encoding DNA is different from the encoding DNA of the first exon of erythropoietin.
- 25 44. The method of Claim 31 wherein the encoding DNA of the exon is the same as the encoding DNA of the first exon of human growth hormone.

-114-

45. The method of Claim 28 further comprising the step of:
(c) maintaining a homologously recombinant cell from step (b) under conditions appropriate for production of a protein.
46. The method of Claim 45 in which the gene whose expression is altered is the erythropoietin gene.
47. Erythropoietin produced by the method of Claim 45.
48. A fusion protein containing amino acids encoded by exons from the DNA construct and amino acids encoded by an endogenous gene produced by the method of Claim 45.
49. A fusion protein of Claim 48 wherein the endogenous gene is erythropoietin.
50. A fusion protein of Claim 49 comprising amino acids 1-3 of human growth hormone and amino acids 6-165 of human erythropoietin.
51. A homologously recombinant cell produced by the method of Claim 28.
52. A homologously recombinant cell produced by the method of Claim 29.
53. A homologously recombinant cell produced by the method of Claim 30.
54. A homologously recombinant cell produced by the method of Claim 31.

-115-

55. A homologously recombinant cell produced by the method of Claim 32.
56. A homologously recombinant cell produced by the method of Claim 33.
- 5 57. A homologously recombinant cell produced by the method of Claim 40.
58. A homologously recombinant cell produced by the method of Claim 41.
- 10 59. A homologously recombinant cell produced by the method of Claim 42.
60. A homologously recombinant cell produced by the method of Claim 44.
- 15 61. A homologously recombinant cell comprising an exogenous regulatory sequence, an exogenous exon and a splice-donor site, operatively linked to the second exon of an endogenous gene.
62. The homologously recombinant cell of Claim 61 wherein the exogenous exon comprises a CAP site.
- 20 63. The homologously recombinant cell of Claim 62 wherein the exogenous exon further comprises the nucleotide sequence ATG.
64. The homologously recombinant cell of Claim 63 wherein the exogenous exon further comprises encoding DNA in-frame with the targeted endogenous gene.

-116-

65. The homologously recombinant cell of Claim 64 wherein the encoding DNA is the same as the encoding DNA of the first exon of the targeted gene.
- 5 66. The homologously recombinant cell of Claim 64 wherein the encoding DNA is different from the encoding DNA of the first exon of the targeted gene.
- 10 67. The homologously recombinant cell of Claim 64 wherein the exogenous regulatory sequence, exogenous exon and splice-donor site are upstream of the coding region of the targeted gene.
68. The homologously recombinant cell of Claim 67 wherein the exogenous regulatory sequence, exogenous exon and splice-donor site are upstream of the endogenous regulatory sequence of the targeted gene.
- 15 69. The homologously recombinant cell of Claim 61 wherein the endogenous regulatory sequence is deleted.
70. The homologously recombinant cell of Claim 69 wherein the first endogenous exon is deleted.
- 20 71. The homologously recombinant cell of Claim 64 wherein the targeted gene encodes a hormone, a cytokine, an antigen, an antibody, an enzyme, a clotting factor, a transport protein, a receptor, a regulatory protein, a structural protein or a transcription factor.
- 25 72. The homologously recombinant cell of Claim 64 wherein the targeted gene encodes a protein selected from the group consisting of erythropoietin, calcitonin, growth hormone, insulin, insulinotropin, insulin-like growth factors, parathyroid hormone, β -inter-

-117-

- feron, γ -interferon, nerve growth factors, FSH β , TGF- β , tumor necrosis factor, glucagon, bone growth factor-2, bone growth factor-7, TSH- β , interleukin 1, interleukin 2, interleukin 3, interleukin 6, interleukin 11, interleukin 12, CSF-granulocyte, CSF-macrophage, CSF-granulocyte/ macrophage, immunoglobulins, catalytic antibodies, protein kinase C, glucocerebrosidase, superoxide dismutase, tissue plasminogen activator, urokinase, antithrombin III, DNase, α -galactosidase, tyrosine hydroxylase, blood clotting factor V, blood clotting factor VII, blood clotting factor VIII, blood clotting factor IX, blood clotting factor X, blood clotting factor XIII, apolipoprotein E or apolipoprotein A-I, globins, low density lipoprotein receptor, IL-2 receptor, IL-2 antagonists, alpha-1 antitrypsin, immune response modifiers, and soluble CD4.
73. The homologously recombinant cell of Claim 61 wherein the cell is a eukaryote.
74. The homologously recombinant cell of Claim 73 wherein the cell is of fungal, plant or animal origin.
75. The homologously recombinant cell of Claim 74 wherein the cell is of vertebrate origin.
76. The homologously recombinant cell of Claim 75 wherein the cell is a primary or secondary mammalian cell.
77. The homologously recombinant cell of Claim 75 wherein the cell is a primary or secondary human cell.
78. The homologously recombinant cell of Claim 75 wherein the cell is an immortalized mammalian cell.

-118-

79. The homologously recombinant cell of Claim 75 wherein the cell is an immortalized human cell.
80. The homologously recombinant cell of Claim 75 wherein the cell is selected from the group consisting of:
5 HT1080 cells, HeLa cells and derivatives of HeLa cells, MCF-7 breast cancer cells, K-562 leukemia cells, KB carcinoma cells, 2780AD ovarian carcinoma cells, Raji cells, Jurkat cells, Namalwa cells, HL-60 cells, Daudi cells, RPMI 8226 cells, U-937 cells,
10 Bowes Melanoma cells, WI-38VA13 subline 2R4 cells, and MOLT-4 cells.
81. The homologously recombinant cell of Claim 80 wherein the targeted gene encodes erythropoietin.
82. The homologously recombinant cell of Claim 81 capable
15 of expressing erythropoietin.
83. The homologously recombinant cell of Claim 82 wherein the encoding DNA is the same as the encoding DNA of the first exon of erythropoietin.
84. The homologously recombinant cell of Claim 81 wherein
20 the encoding DNA is different from the encoding DNA of the first exon of erythropoietin.
85. The homologously recombinant cell of Claim 84 wherein the encoding DNA is the same as the encoding DNA of the first exon of human growth hormone.
- 25 86. The homologously recombinant cell of Claim 61 capable of expressing a fusion protein comprising amino acids encoded by the exogenous exon and amino acids encoded by the endogenous gene.

-119-

87. A fusion protein of Claim 86 wherein the endogenous gene is erythropoietin.
88. A fusion protein of Claim 87 comprising amino acids 1-3 of human growth hormone and amino acids 6-165 of human erythropoietin.
89. The homologously recombinant cell of Claim 66 wherein the regulatory sequence is a promoter, an enhancer, a scaffold-attachment region or a transcription factor binding site.
90. The homologously recombinant cell of Claim 89 wherein the exogenous regulatory sequence is a promoter.
91. The homologously recombinant cell of Claim 89 wherein the exogenous regulatory sequence is a regulatory sequence of the mouse metallothionein-I gene, a regulatory sequence of an SV-40 gene, a regulatory sequence of a cytomegalovirus gene, a regulatory sequence of a collagen gene, a regulatory sequence of an actin gene, a regulatory sequence of an immunoglobulin gene, a regulatory sequence of the HMG-CoA reductase gene or a regulatory sequence of the EF-1 α gene.
92. A method of altering the expression of a gene in a cell, comprising the steps of:
- (a) transfecting a cell with a DNA construct, the DNA construct comprising:
- (i) a targeting sequence;
- (ii) a regulatory sequence;
- (iii) an exon; and
- (iv) an unpaired splice-donor site, thereby producing a transfected cell;

-120-

- (b) maintaining the transfected cell under conditions appropriate for homologous recombination, thereby producing a homologously recombinant cell; and
 - 5 (c) maintaining the homologously recombinant cell under conditions appropriate for expression of the gene.
- 93. The method of Claim 92 wherein the exon comprises the nucleotide sequence ATG.
- 10 94. The method of Claim 92 wherein the exon further comprises a CAP site.
- 95. The method of Claim 94 wherein the exon further comprises encoding DNA which is in-frame with the targeted gene.
- 15 96. The method of Claim 95 wherein the encoding DNA is the same as the encoding DNA of the first exon of the targeted gene.
- 97. The method of Claim 96 wherein the targeted gene is the erythropoietin gene.
- 20 98. The method of Claim 96 wherein the encoding DNA is different from the encoding DNA of the first exon of the targeted gene.
- 99. The method of Claim 98 wherein the targeted gene is the erythropoietin gene.
- 25 100. The method of Claim 98 wherein the targeting sequence is homologous to a sequence within the targeted gene.

-121-

101. The method of Claim 98 wherein the targeting sequence is homologous to a sequence upstream of the coding region of the targeted gene.
- 5 102. The method of Claim 98 wherein the targeting sequence is homologous to a sequence upstream of the endogenous regulatory sequence for the targeted gene.
103. The method of Claim 98 wherein the construct further comprises a second targeting sequence homologous to a sequence within the targeted gene.
- 10 104. The method of Claim 98 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the coding region of the targeted gene.
- 15 105. The method of Claim 98 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the endogenous regulatory sequence for the targeted gene.
106. The method of Claim 92 further comprising the step of:
- 20 (c) maintaining a homologously recombinant cell under conditions appropriate for production of a protein.
107. The method of Claim 106 in which the gene whose expression is altered is the erythropoietin gene.
- 25 108. Erythropoietin produced by the method of Claim 107.
109. A fusion protein produced by the method of Claim 106.

-122-

110. A fusion protein of Claim 109 wherein the endogenous gene is erythropoietin.
111. A fusion protein of Claim 110 comprising amino acids 1-3 of human growth hormone and amino acids 6-165 of human erythropoietin.
112. A method of making a protein by altering the expression of a gene in a cell, comprising the steps of:
- (a) transfecting a cell with a DNA construct, the DNA construct comprising:
 - (i) a targeting sequence;
 - (ii) a regulatory sequence;
 - (iii) an exon; and
 - (iv) an unpaired splice-donor site, thereby producing a transfected cell;
 - (b) maintaining the transfected cell under conditions appropriate for homologous recombination, thereby producing a homologously recombinant cell; and
 - (c) maintaining the homologously recombinant cell under conditions appropriate for production of the protein.
113. The method of Claim 112 wherein the exon comprises a CAP site.
114. The method of Claim 113 wherein the exon comprises the nucleotide sequence ATG.
115. The method of Claim 114 wherein the exon further comprises encoding DNA which is in-frame with the targeted endogenous gene.

-123-

116. The method of Claim 115 wherein the encoding DNA is the same as the encoding DNA of the first exon of the targeted gene.
- 5 117. The method of Claim 116 wherein the encoding DNA is different from the encoding DNA of the first exon of the targeted gene.
118. The method of Claim 117 wherein the targeting sequence is homologous to a sequence within the targeted gene.
- 10 119. The method of Claim 117 wherein the targeting sequence is homologous to a sequence upstream of the coding region of the targeted gene.
120. The method of Claim 117 wherein the targeting sequence is homologous to a sequence upstream of the endogenous regulatory sequence for the targeted gene.
- 15 121. The method of Claim 117 wherein the construct further comprises a second targeting sequence homologous to a sequence within the targeted gene.
122. The method of Claim 117 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the coding region of the targeted gene.
- 20 123. The method of Claim 117 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the endogenous regulatory sequence for the targeted gene.
- 25 124. An erythropoietin produced by the method of Claim 112.

-124-

125. The erythropoietin of Claim 124 wherein the cell is of human origin.
126. A protein produced by the method of Claim 112.
127. The protein of Claim 126 which is a fusion protein.
- 5 128. The fusion protein of Claim 127 wherein the endogenous gene is the erythropoietin gene.
129. The fusion protein of Claim 128 comprising amino acids 1-3 of human growth hormone and amino acids 6-165 of human erythropoietin.
- 10 130. The DNA plasmid pREPO18.
131. A DNA construct capable of altering the expression of a targeted gene when inserted into the chromosomal DNA of a cell, comprising:
- 15 (a) a targeting sequence;
- (b) a regulatory sequence;
- (c) an exon;
- (d) a splice-donor site;
- (e) an intron; and
- (f) a splice-acceptor site.
- 20 132. The DNA construct of Claim 131 wherein the targeting sequence is homologous to a sequence within the targeted gene.
133. The DNA construct of Claim 131 wherein the targeting sequence is homologous to a sequence upstream of the coding region of the targeted gene.
- 25

-125-

134. The DNA construct of Claim 131 wherein the targeting sequence is homologous to a sequence upstream of the endogenous regulatory sequence of the targeted gene.
- 5 135. The DNA construct of Claim 131 wherein the construct further comprises a second targeting sequence homologous to a sequence within the targeted gene.
- 10 136. The DNA construct of Claim 131 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the coding region of the targeted gene.
137. The DNA construct of Claim 131 wherein the construct further comprises a second targeting sequence homologous to a sequence upstream of the endogenous regulatory sequence of the targeted gene.
- 15 138. A homologously recombinant cell comprising a regulatory sequence, an exon, a splice-donor site, an intron and a splice-acceptor site introduced by homologous recombination upstream of the coding region of a targeted gene.
- 20 139. The homologously recombinant cell of Claim 138 wherein the targeted gene is the α -interferon gene.
140. The homologously recombinant cell of Claim 138 wherein the targeted gene is the erythropoietin gene.
- 25 141. A homologously recombinant cell comprising the dhfr gene, the neo gene, the CMV promoter, hGH exon 1 and an unpaired splice-donor site targeted to a position upstream of the endogenous erythropoietin regulatory region.

-126-

142. The homologously recombinant cell of Claim 141 produced by the integration of DNA from pREPO18.
143. A method of making a homologously recombinant cell wherein the expression of a targeted gene is altered, comprising the steps of:
- 5 (a) transfecting a cell with a DNA construct, the construct comprising:
- 10 (i) a targeting sequence;
- (ii) a regulatory sequence;
- (iii) an exon;
- (iv) a splice-donor site;
- (v) an intron; and
- (vi) a splice-acceptor site;
- 15 wherein the targeting sequence directs the integration of elements (b)-(f) upstream such that they are operatively linked to the first exon of a targeted gene; and
- (b) maintaining the transfected cell under conditions appropriate for homologous recombination.
- 20 144. A homologously recombinant cell produced by the method of Claim 143.
145. A method of altering the expression of a gene in a cell, comprising the steps of:
- 25 (a) transfecting a cell with a DNA construct, the construct comprising:
- (i) a targeting sequence;
- (ii) a regulatory sequence;
- (iii) an exon;
- (iv) a splice-donor site;
- 30 (v) an intron; and
- (vi) a splice-acceptor site;

-127-

wherein the targeting sequence directs the integration of elements (b)-(f) upstream such that they are operatively linked to the first exon of a targeted gene;

- 5 (b) maintaining the transfected cell under conditions appropriate for homologous recombination; and
- (c) maintaining the homologously recombinant cell under conditions appropriate for expression of
- 10 the gene.

146. A method of making a protein by altering the expression of a gene in a cell, comprising the steps of:

- (a) transfecting a cell with a DNA construct, the construct comprising:
- 15 (i) a targeting sequence;
- (ii) a regulatory sequence;
- (iii) an exon;
- (iv) a splice-donor site;
- (v) an intron; and
- 20 (vi) a splice-acceptor site;
- wherein the targeting sequence directs the integration of elements (b)-(f) upstream such that they are operatively linked to the first exon of a targeted gene;
- 25 (b) maintaining the transfected cell under conditions appropriate for homologous recombination; and
- (c) maintaining the homologously recombinant cell under conditions appropriate for expression of
- 30 the protein.

147. The method of Claim 146 wherein the targeted gene is the α -interferon gene or the erythropoietin gene.

-128-

148. DNA sequences located between about 5 kilobases and 30 kilobases upstream of the ATG of the erythropoietin gene.
- 5 149. A method for targeting the erythropoietin gene in a mammalian cell comprising transfecting the cell with a construct comprising a DNA sequence homologous to a sequence upstream of the sequence ATG of the erythropoietin gene.
- 10 150. The method of Claim 149 wherein the construct comprising a DNA sequence homologous to a sequence located between about 5 kilobases and 30 kilobases upstream of the sequence ATG of the erythropoietin gene.
- 15 151. The method of Claim 150 wherein the mammalian cell is a human cell.
152. A method for targeting the erythropoietin gene in a mammalian cell comprising transfecting the cell with a construct comprising a DNA sequence homologous to a sequence within the erythropoietin gene.
- 20 153. The method of Claim 152 wherein the mammalian cell is a human cell.

1/15




FIGURE 1

SUBSTITUTE SHEET (RULE 26)

2/15




FIGURE 2

SUBSTITUTE SHEET (RULE 26)

3/15




FIG. 3

SUBSTITUTE SHEET (RULE 26)

4/15





FIGURE 4

SUBSTITUTE SHEET (RULE 26)

5/15

**FIG. 5****SUBSTITUTE SHEET (RULE 26)**

6/15




FIGURE 6

SUBSTITUTE SHEET (RULE 26)

7/15

EPO Production in Targeted HT1080
Cell Line 52 Under MTX Selection




Figure 7

SUBSTITUTE SHEET (RULE 26)

8/15




FIGURE 8

SUBSTITUTE SHEET (RULE 26)

9/15





FIGURE 9A

10/15

FIG. 9B

SUBSTITUTE SHEET (RULE 26)

11/15




FIGURE 10

SUBSTITUTE SHEET (RULE 26)

12/15





FIGURE 11

SUBSTITUTE SHEET (RULE 26)

13/15



SUBSTITUTE SHEET (RULE 26)




FIGURE 13

15/15




Figure 14

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

 Interns 1 Application No
 PCT/US 95/06045

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/67 C12N15/90 C12N15/85 C12N15/62 C12N5/06 C07K14/505 C07K14/61 A61K48/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 05784 (THE UNITED STATES OF AMERICA) 17 March 1994	1,4,7, 21,28, 31,34, 40,45, 51-57, 92,95, 100,106, 112,115, 118,131, 132,138 143-146
X	see page 4, line 3 - line 9 see page 4, line 21 - page 5, line 26 see page 13, line 3 - page 15, line 13 see page 21, line 25 - page 22, line 30 see page 23, line 20 - line 31 see page 24, line 14 - line 32; examples 3-8 --- -/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "A" document member of the same patent family		
Date of the actual completion of the international search 28 July 1995		Date of mailing of the international search report 1 0. 08. 95
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer Montero Lopez, B

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/US 95/06045

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	WO,A,91 09955 (APPLIED RESEARCH SYSTEMS, ARS HOLDING MV) 11 July 1991 cited in the application see page 1, line 6 - line 26 see page 6, line 15 - page 8, line 27 see page 16, line 4 - page 18, line 5 see page 18, line 22 - page 19, line 28 see page 21, line 14 - page 22, line 6 see page 23, line 4 - page 28, line 21 ---	51-56, 126,144 1-46, 61-107, 112-123, 130-143, 145-147
X A	WO,A,91 06667 (CELL GENESYS, INC.) 16 May 1991 cited in the application see page 3, line 12 - line 26 see page 4, line 6 - page 6, line 38 see page 8, line 29 - page 9, line 9 see page 12, line 33 - line 34 see page 15, line 4 - page 16, line 38 ---	47, 51-60, 108, 124-126, 144, 148-153 1-46, 61-107, 112-123, 130-143, 145-147
P,X	WO,A,94 12650 (TRANSKARYOTIC THERAPIES, INC.) 9 June 1994 see page 1, line 29 - page 4, line 16 see page 5, line 27 - page 6, line 4 see page 9, line 3 - page 14, line 10 see page 16, line 20 - page 23, line 20 see page 30, line 1 - page 31, line 2; examples 1-6 -----	1-153

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat' Application No
PCT/US 95/06045

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9405784	17-03-94	AU-B- 4839493	29-03-94
WO-A-9109955	11-07-91	AU-B- 645294	13-01-94
		AU-A- 7183691	24-07-91
		CA-A- 2071989	23-06-91
		EP-A- 0505500	30-09-92
		JP-T- 5504682	22-07-93
		QA-A- 9594	30-04-93
		US-A- 5272071	21-12-93
WO-A-9106667	16-05-91	AU-B- 635844	01-04-93
		AU-A- 7740791	31-05-91
		EP-A- 0452484	23-10-91
		JP-T- 4505104	10-09-92
		WO-A- 9106666	16-05-91
WO-A-9412650	09-06-94	AU-B- 5736294	22-06-94